For research purposes only, not for human use

Product Data Sheet

Anti-RPL28 Antibody

Catalog #	Source	Reactivity	Applications
CPA2022	Rabbit	H, M, R, Mk	WB, IH

Description	Rabbit polyclonal antibody to RPL28
Immunogen	KLH-conjugated synthetic peptide encompassing a sequence within the center region of human RPL28. The exact sequence is proprietary.
Purification	The antibody was purified by immunogen affinity chromatography.
Specificity	Recognizes endogenous levels of RPL28 protein.
Clonality	Polyclonal
Conjugation	
Form	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.01% sodium azide.
Dilution	WB (1/500 - 1/1000), IH (1/50 - 1/100)
Gene Symbol	RPL28
Alternative Names	60S ribosomal protein L28
Entrez Gene	6158 (Human)
SwissProt	P46779 (Human)
Storage/Stability	Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid freeze/thaw cycles.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference
Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB www.cohesionbio.com
ORDER order@cohesionbio.com
SUPPORT techsupport@cohesionbio.com
CUSTOM custom@cohesionbio.com

For research purposes only, not for human use

Product Data Sheet



Western blot analysis of RPL28 expression in HEK293T (A), Hela (B), H446 (C), mouse liver (D), mouse kidney (E), rat liver (F), rat kidney (G) whole cell lysates. (Predicted band size: 15 kD; Observed band size: 16 kD)



Immunohistochemical analysis of RPL28 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB www.cohesionbio.com
ORDER order@cohesionbio.com
SUPPORT techsupport@cohesionbio.com
CUSTOM custom@cohesionbio.com