

Product Data Sheet

Anti-Tetraspanin 8 Antibody

Catalog #	Source	Reactivity	Applications		
CPA2172	Rabbit	H, M, R	WB, IH, IF/IC		
Description	R	abbit polyclonal antibody t	o Tetraspanin 8		
Immunogen	K	LH-conjugated synthetic pe	ptide encompassing a sequence within the center		
	re	egion of human Tetraspanir	8. The exact sequence is proprietary.		
Purification	Т	he antibody was purified by	/ immunogen affinity chromatography.		
Specificity	R	ecognizes endogenous leve	ls of Tetraspanin 8 protein.		
Clonality	lonality Polyclonal				
Conjugation					
Form		Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,			
	а	nd 0.01% sodium azide.			
Dilution	V	VB (1/500 - 1/1000), IH (1/50	- 1/100), IF/IC (1/50 - 1/200)		
Gene Symbol	Т	SPAN8			
Alternative Na	ames T	M4SF3; Tetraspanin-8; Tspa	n-8; Transmembrane 4 superfamily member 3;		
	Т	umor-associated antigen CO	0-029		
Entrez Gene	7	'103 (Human)			
SwissProt	Р	19075 (Human)			
Storage/Stabi	lity S	hipped at 4°C. Upon deliver	y aliquot and store at -20°C for one year. Avoid		
	fr	reeze/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish


COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com




For research purposes only, not for human use

Product Data Sheet



Western blot analysis of Tetraspanin 8 expression in HEK293T (A), Hela (B), HGC27 (C), mouse testis (D), rat testis (E) whole cell lysates. (Predicted band size: 26 kD; Observed band size: 35 kD)



Immunohistochemical analysis of Tetraspanin 8 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of Tetraspanin 8 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com