

Product Data Sheet

Anti-Collagen 5 alpha 3 Antibody

Catalog #	Source	Reactivity	Applications		
CPA2654	Rabbit	H, M, Mk	WB, IH, IF/IC		
Description	F	Rabbit polyclonal antibody to Collagen 5 alpha 3			
Immunogen	ķ	KLH-conjugated synthetic peptide encompassing a sequence within the N-term			
	r	region of human Collagen 5 alpha 3. The exact sequence is proprietary.			
Purification	T	The antibody was purified b	y immunogen affinity chromatography.		
Specificity	F	Recognizes endogenous leve	els of Collagen 5 alpha 3 protein.		
Clonality	F	Polyclonal			
Conjugation					
Form		Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,			
	a	and 0.01% sodium azide.			
Dilution	١	WB (1/500 - 1/1000), IH (1/50	- 1/100), IF/IC (1/50 - 1/200)		
Gene Symbol	C	COL5A3			
Alternative Na	ames (Collagen alpha-3(V) chain			
Entrez Gene	5	50509 (Human)			
SwissProt	F	25940 (Human)			
Storage/Stabi	lity S	Shipped at 4°C. Upon delive	ry aliquot and store at -20°C for one year. Avoid		
	f	reeze/thaw cycles.			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish


COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com




For research purposes only, not for human use

Product Data Sheet



Western blot analysis of Collagen 5 alpha 3 expression in CT26 (A), MCF7 (B), Hela (C) whole cell lysates. (Predicted band size: 172 kD; Observed band size: 180; 120 kD)



Immunohistochemical analysis of Collagen 5 alpha 3 staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Immunofluorescent analysis of Collagen 5 alpha 3 staining in HuvEc cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com