

Product Data Sheet

Anti-p67 Antibody

Catalog # Source Reactivity Applications

CPA4842 Rabbit H, M, R, B WB, IH, IF/IC

Description Rabbit polyclonal antibody to p67

Immunogen KLH-conjugated synthetic peptide encompassing a sequence within the center

region of human p67. The exact sequence is proprietary.

Purification The antibody was purified by immunogen affinity chromatography.

Specificity Recognizes endogenous levels of p67 protein.

Clonality Polyclonal

Conjugation

Form Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,

and 0.01% sodium azide.

Dilution WB (1/500 - 1/1000), IH (1/100 - 1/200), IF/IC (1/100 - 1/500)

Gene Symbol STXBP1

Alternative Names UNC18A; Syntaxin-binding protein 1; MUNC18-1; N-Sec1; Protein unc-18 homolog 1;

Unc18-1; Protein unc-18 homolog A; Unc-18A; p67

Entrez Gene 6812 (Human); 20910 (Mouse); 25558 (Rat)

SwissProt P61764 (Human); O08599 (Mouse); P61765 (Rat)

Storage/Stability Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid


freeze/thaw cycles.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish


COHESION BIOSCIENCES LIMITED




Product Data Sheet



Western blot analysis of p67 expression in U87MG (A), C6 (B), SHSY5Y (C), K562 (D), Raw264.7 (E), mouse brain (F), rat brain (G) whole cell lysates. (Predicted band size: 67 kD; Observed band size: 70 kD)



Immunohistochemical analysis of p67 staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of p67 staining in Jurkat cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED