

Product Data Sheet

Anti-RAB5C Antibody

Catalog #	Source	e Reactivity	Applications
CPA4934	Rabbit	H, M, R, B, D	WB, IH
Description		Rabbit polyclonal antibody to	RAB5C
Immunogen		KLH-conjugated synthetic per	tide encompassing a sequence within the center
		region of human RAB5C. The	exact sequence is proprietary.
Purification		The antibody was purified by	immunogen affinity chromatography.
Specificity		Recognizes endogenous level	s of RAB5C protein.
Clonality		Polyclonal	
Conjugation			
Form		Liquid in 0.42% Potassium ph	osphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,
		and 0.01% sodium azide.	
Dilution		WB (1/500 - 1/1000), IH (1/50 -	1/100)
Gene Symbol		RAB5C	
Alternative N	ames	RABL; Ras-related protein Rab	o-5C; L1880; RAB5L
Entrez Gene		5878 (Human); 19345 (Mouse	e)
SwissProt		P51148 (Human); P35278 (M	ouse)
Storage/Stabi	lity	Shipped at 4°C. Upon delivery	aliquot and store at -20°C for one year. Avoid
		freeze/thaw cycles.	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED


WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



KDa A B C D 70 45 35 25 20 15 For research purposes only, not for human use

Product Data Sheet

Western blot analysis of RAB5C expression in Hela (A), HGC27 (B), mouse lung (C), rat lung (D) whole cell lysates. (Predicted band size: 23 kD; Observed band size: 24 kD)



Immunohistochemical analysis of RAB5C staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com