

Product Data Sheet

Anti-Alpha-1D Adrenergic Receptor Antibody

Catalog #	Source	Reactivity	Applications	
CPA5319	Rabbit	Н	WB, IH, IF/IC	
Description	Rabl	oit polyclonal antibody	y to Alpha-1D Adrenergic Receptor	
Immunogen	KLH-	conjugated synthetic	peptide encompassing a sequence within the C-term	
	regio	on of human Alpha-1D	Adrenergic Receptor. The exact sequence is proprietary.	
Purification	The	antibody was purified	by immunogen affinity chromatography.	
Specificity	Reco	gnizes endogenous le	vels of Alpha-1D Adrenergic Receptor protein.	
Clonality	Poly	clonal		
Conjugation				
Form	Liqu	id in 0.42% Potassium	phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,	
	and	0.01% sodium azide.		
Dilution	WB	(1/500 - 1/2000), IH (1/5	50 - 1/200), IF/IC (1/50 - 1/100)	
Gene Symbol	ADR	A1D		
Alternative Na	mes ADR	A1A; Alpha-1D adrene	ergic receptor; Alpha-1A adrenergic receptor; Alpha-1D	
	adre	noreceptor; Alpha-1D	adrenoceptor; Alpha-adrenergic receptor 1a	
Entrez Gene	146	(Human); 29413 (Rat)		
SwissProt	P251	100 (Human); P97714	(Mouse); P23944 (Rat)	
Storage/Stabil	ity Ship	ped at 4°C. Upon deliv	very aliquot and store at -20°C for one year. Avoid	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish


freeze/thaw cycles.

COHESION BIOSCIENCES LIMITED


WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com




Product Data Sheet



Western blot analysis of Alpha-1D Adrenergic Receptor expression in H9C2 (A), Raw264.7 (B), U87MG (C) whole cell lysates. (Predicted band size: 60 kD; Observed band size: 60 kD)



Immunohistochemical analysis of Alpha-1D Adrenergic Receptor staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of Alpha-1D Adrenergic Receptor staining in MCF7 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark. DAPI was used to stain the cell nuclei (blue).

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED