

Product Data Sheet

Anti-OATP2 Antibody

Catalog #	Source	e Reactivity	Applications	
CPA5637	Rabbit	t H	WB, IH	
Description		Rabbit polyclonal antibody	to OATP2	
Immunogen		KLH-conjugated synthetic peptide encompassing a sequence within the center		
		region of human OATP2. Th	ne exact sequence is proprietary.	
Purification		The antibody was purified	by immunogen affinity chromatography.	
Specificity		Recognizes endogenous lev	vels of OATP2 protein.	
Clonality		Polyclonal		
Conjugation				
Form		Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,		
		and 0.01% sodium azide.		
Dilution		WB (1/500 - 1/1000), IH (1/5	0 - 1/100)	
Gene Symbol		SLCO1B1		
Alternative Names		LST1; OATP1B1; OATP2; OATPC; SLC21A6; Solute carrier organic anion transporter		
		family member 1B1; Liver-	specific organic anion transporter 1; LST-1; OATP-C;	
		Sodium-independent organ	nic anion-transporting polypeptide 2; OATP-2; Solute	
		carrier family 21 member 6		
Entrez Gene		10599 (Human)		
SwissProt		Q9Y6L6 (Human)		
Storage/Stabi	lity	Shipped at 4°C. Upon deliv	ery aliquot and store at -20°C for one year. Avoid	
		freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish


COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com


Cohesion

For research purposes only, not for human use

Product Data Sheet



Western blot analysis of OATP2 expression in LO2 (A) whole cell lysates. (Predicted band size: 76 kD; Observed band size: 90 kD)



Immunohistochemical analysis of OATP2 staining in human tonsil cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com