

Product Data Sheet

Anti-HMGN2 (Phospho-S29) Antibody

Catalog # Source Reactivity Applications

CPA5882 Rabbit H, M, R, B, C, D, P WB, IH

Description Rabbit polyclonal antibody to HMGN2 (Phospho-S29)

Immunogen KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding

S29 of human HMGN2 protein. The exact sequence is proprietary.

Purification The antibody was purified by immunogen affinity chromatography.

Specificity Recognizes endogenous levels of HMGN2 protein only when phosphorylated at S29.

Clonality Polyclonal

Conjugation

Form Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,

and 0.01% sodium azide.

Dilution WB (1/500 - 1/1000), IH (1/50 - 1/200)

Gene Symbol HMGN2

Alternative Names HMG17; Non-histone chromosomal protein HMG-17; High mobility group

nucleosome-binding domain-containing protein 2

Entrez Gene 3151 (Human); 15331 (Mouse)

SwissProt P05204 (Human); P09602 (Mouse); P18437 (Rat)

Storage/Stability Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid

freeze/thaw cycles.


Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED


WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com



Product Data Sheet



Western blot analysis of HMGN2 (Phospho-S29) expression in rat lung (A) whole cell lysates. (Predicted band size: 9 kD; Observed band size: 17 kD)



Immunohistochemical analysis of HMGN2 (Phospho-S29) staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED