

Product Data Sheet

Anti-WTAP Antibody

Catalog #	Source	Reactivity	Applications	
CPA6128	Rabbit	-	WB, IH	
Description		Rabbit polyclonal antibody t	o WTAP	
Immunogen		KLH-conjugated synthetic pe	ptide encompassing a sequence within the C-term	
		region of human WTAP. The	exact sequence is proprietary.	
Purification		The antibody was purified by	<i>immunogen affinity chromatography.</i>	
Specificity		Recognizes endogenous leve	ls of WTAP protein.	
Clonality		Polyclonal		
Conjugation				
Form		Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,		
		and 0.01% sodium azide.		
Dilution		WB (1/500 - 1/1000), IH (1/50	- 1/200)	
Gene Symbol		WTAP		
Alternative Na	ames	KIAA0105; Pre-mRNA-splicin	g regulator WTAP; Female-lethal(2)D homolog; hFL(2)D;	
		WT1-associated protein; Wil	ms tumor 1-associating protein	
Entrez Gene		9589 (Human); 60532 (Mou	se)	
SwissProt		Q15007 (Human); Q9ER69 (I	Mouse)	
Storage/Stabi	lity	Shipped at 4°C. Upon deliver	y aliquot and store at -20°C for one year. Avoid	
		freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED


WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com

Cohesion

For research purposes only, not for human use

Product Data Sheet

kDa A B C 180 130 95 72 55 43 Western blot analysis of WTAP expression in MEF (A), EC9706 (B), Panc1 (C) whole cell lysates. (Predicted band size: 44 kD; Observed band size: 55 kD)



Immunohistochemical analysis of WTAP staining in human brain formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com