

Product Data Sheet

Anti-BLNK (Phospho-Y96) Antibody

Catalog # Source Reactivity Applications

CPA7139 Rabbit H, Mk WB, IH

Description Rabbit polyclonal antibody to BLNK (Phospho-Y96)

Immunogen KLH-conjugated synthetic phosphopeptide corresponding to residues surrounding

Y96 of human BLNK protein. The exact sequence is proprietary.

Purification The antibody was purified by immunogen affinity chromatography.

Specificity Recognizes endogenous levels of BLNK protein only when phosphorylated at Y96.

Clonality Polyclonal

Conjugation

Form Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,

and 0.01% sodium azide.

Dilution WB (1/500 - 1/1000), IH (1/100 - 1/200)

Gene Symbol BLNK

Alternative Names BASH; SLP65; B-cell linker protein; B-cell adapter containing a SH2 domain protein;

B-cell adapter containing a Src homology 2 domain protein; Cytoplasmic adapter

protein; Src homology 2 domain-containing leukocyte protein of 65 kDa; SLP-65

Entrez Gene 29760 (Human)

SwissProt Q8WV28 (Human)

Storage/Stability Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid

freeze/thaw cycles.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish


COHESION BIOSCIENCES LIMITED

WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com






Product Data Sheet



Western blot analysis of BLNK (Phospho-Y96) expression in HT29 (A), HEK293T (B) whole cell lysates. (Predicted band size: 50 kD; Observed band size: 70 kD)



Immunohistochemical analysis of BLNK (Phospho-Y96) staining in human lymph node formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED