

Product Data Sheet

Anti-Histone H2A (Acetyl-K5) Antibody

Catalog #	Source	Reactivity	Applications	
CPA9315	Rabbit	H, M, R	WB, IH	
Description		Rabbit polyclonal antibody t	o Histone H2A (Acetyl-K5)	
Immunogen	I	KLH-conjugated synthetic ac	etylated peptide corresponding to residues surrounding	
	I	K5 of human Histone H2A pr	otein. The exact sequence is proprietary.	
Purification	-	The antibody was purified by	immunogen affinity chromatography.	
Specificity	I	Recognizes endogenous leve	ls of Histone H2A protein only when acetylated at K5.	
Clonality	I	Polyclonal		
Conjugation				
Form	I	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,		
	i	and 0.01% sodium azide.		
Dilution		WB (1/1000 - 1/2000), IH (1/2	00 - 1/500)	
Gene Symbol	I	HIST1H2AG; HIST1H2AI; HIST	1H2AK; HIST1H2AL; HIST1H2AM; HIST2H2AA3;	
	I	HIST2H2AA4; HIST3H2A		
Alternative N	ames	HIST1H1A; H1F1; Histone H1	.1; Histone H1a; HIST1H1C; H1F2; Histone H1.2; Histone	
	I	H1c; Histone H1d; Histone H	1s-1; HIST1H1D; H1F3; Histone H1.3; Histone H1c;	
	I	Histone H1s-2; HIST1H1E; H2	.F4; Histone H1.4; Histone H1b; Histone H1s-4	
Entrez Gene	;	8329, 8330, 8332, 8336, 896	9, 723790, 8337, 92815 (Human); 319164, 15267,	
	:	319162 (Mouse); 365877, 64	646 (Rat)	
SwissProt	I	P0C0S8, Q6FI13, Q7L7L0 (Hu	man); P22752, Q6GSS7, Q8BFU2 (Mouse); P02262,	
	I	POCC09, Q4FZT6 (Rat)		
Storage/Stabi	lity	Shipped at 4°C. Upon deliver	y aliquot and store at -20°C for one year. Avoid	
	ł	freeze/thaw cycles.		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish


COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com




For research purposes only, not for human use

Product Data Sheet



Western blot analysis of Histone H2A (Acetyl-K5) expression in Hela TSA-treated (A) whole cell lysates. (Predicted band size: 14 kD; Observed band size: 14 kD)



Immunohistochemical analysis of Histone H2A (Acetyl-K5) staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com