

Product Data Sheet

Anti-Cystinosin Antibody


Catalog #	Source	Reactivity	Applications
CQA3236	Rabbit	H, M, R	WB, IF/IC
Description		Rabbit polyclonal antibody t	o Cystinosin
Immunogen		Recombinant full length pro	tein of human Cystinosin
Purification The antibody was purified by immunogen affinity chromatography.			
Specificity		Recognizes endogenous leve	els of Cystinosin protein.
Clonality		Polyclonal	
Conjugation			
Form	Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol		
		and 0.01% sodium azide.	
Dilution		WB (1/500 - 1/2000), IF/IC (1/	50 - 1/200)
Gene Symbol		CTNS	
Alternative Na	ames	Cystinosin	
Entrez Gene		1497 (Human); 83429 (Mou	se)
SwissProt		O60931 (Human); P57757 (N	/louse)
Storage/Stabi	lity	Shipped at 4°C. Upon delive	ry aliquot and store at -20°C for one year. Avoid
		freeze/thaw cycles.	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com






For research purposes only, not for human use

Product Data Sheet

Western blot analysis of Cystinosin expression in Hela (A), HepG2 (B), mouse kidney (C), rat kidney (D) whole cell lysates. (Predicted band size: 41; 45 kD; Observed band size: 42 kD)



Immunofluorescent analysis of Cystinosin staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a humidified chamber. Cells were washed with PBST and incubated with a DyLight 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com