

## **Product Data Sheet**

## Anti-PI3K p85 gamma Antibody

Catalog # Source Reactivity Applications

CQA5480 Rabbit H, M, R WB, IF/IC

**Description** Rabbit polyclonal antibody to PI3K p85 gamma

Immunogen Recombinant fusion protein of human PI3K p85 gamma. The exact sequence is

proprietary.

**Purification** The antibody was purified by immunogen affinity chromatography.

**Specificity** Recognizes endogenous levels of PI3K p85 gamma protein

**Clonality** Polyclonal

Conjugation

Form Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol,

and 0.01% sodium azide.

**Dilution** WB (1/500 - 1/1000), IF/IC (1/50 - 1/200)

Gene Symbol PIK3R3

Alternative Names Phosphatidylinositol 3-kinase regulatory subunit gamma; PI3-kinase regulatory

subunit gamma; PI3K regulatory subunit gamma; PtdIns-3-kinase regulatory subunit

gamma; Phosphatidylinositol 3-kinase 55 kDa regulatory subunit gamma; PI3-kinase

subunit p55-gamma; PtdIns-3-kinase regulatory subunit p55-gamma; p55PIK

Entrez Gene 8503 (Human); 18710 (Mouse); 60664 (Rat)

SwissProt Q92569 (Human); Q64143 (Mouse); Q63789 (Rat)

Storage/Stability Shipped at 4°C. Upon delivery aliquot and store at -20°C for one year. Avoid

freeze/thaw cycles.


Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## COHESION BIOSCIENCES LIMITED


WEB ORDER SUPPORT CUSTOM
www.cohesionbio.com order@cohesionbio.com techsupport@cohesionbio.com custom@cohesionbio.com



## **Product Data Sheet**



Western blot analysis of PI3K p85 gamma expression in mouse brain (A) whole cell lysates. (Predicted band size: 43; 47; 54 kD; Observed band size: 55 kD)



Immunofluorescent analysis of PI3K p85 gamma staining in NIH3T3 cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a Alexa Fluor 594-conjugated secondary antibody (red) in PBS at room temperature in the dark.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish