

Product Data Sheet

Anti-CLSTN1 Antibody

Catalog #	Source	Reactivity	Applications
CQA8181	Rabbit	H, M, Ha	WB, IH, IF/IC
Description	Rab	bit monoclonal antibod	y to CLSTN1
Immunogen	A sy	ynthetic peptide of hum	an CLSTN1
Purification	The	antibody was purified	oy immunogen affinity chromatography.
Specificity	Rec	ognizes endogenous lev	els of CLSTN1 protein.
Clonality	Мо	noclonal	
Conjugation			
Form	Liqu	uid in 50mM Tris-Glycin	e (pH 7.4), 0.15M NaCl, 50% Glycerol, 0.01% Sodium azide
	and	1 0.05% BSA.	
Dilution	WB	(1/500 - 1/1000), IH (1/5	0 - 1/100), IF/IC (1/50 - 1/100)
Gene Symbol	CLS	TN1	
Alternative Na	imes CS1	.; KIAA0911; Calsyntenir	n-1; Alcadein-alpha; Alc-alpha; Alzheimer-related
	cad	herin-like protein; Non-	classical cadherin XB31alpha
Entrez Gene	228	883 (Human); 65945 (Mo	ouse)
SwissProt	094	4985 (Human); Q9EPL2	(Mouse)
Storage/Stabil	<mark>ity</mark> Shij	pped at 4°C. Upon delive	ery aliquot and store at -20°C for one year. Avoid
	free	eze/thaw cycles.	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED


WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com

Coherion


For research purposes only, not for human use

Product Data Sheet

Western blot analysis of CLSTN1 expression in mouse brain (A), Hela (B), CHOK1 (C) whole cell lysates. (Predicted band size: 110 kD; Observed band size: 115 kD)



Immunohistochemical analysis of CLSTN1 staining in human colon cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.



Immunofluorescent analysis of CLSTN1 staining in HeLa cells. Formalin-fixed cells were permeabilized with 0.1% Triton X-100 in TBS for 5-10 minutes and blocked with 3% BSA-PBS for 30 minutes at room temperature. Cells were probed with the primary antibody in 3% BSA-PBS and incubated overnight at 4 °C in a hidified chamber. Cells were washed with PBST and incubated with a AF488-conjugated secondary antibody (green) in PBS at room temperature in the dark.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com