

# **Product Data Sheet**

## **RUNX2 siRNA (Human)**

rce	Reactivity	Applications		
thetic	Н	RNAi		
siRNA	to inhibit RUNX2 ex	pression using RNA interference		
RUNX	RUNX2 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
knock	down gene expressi	on.		
Lyoph	ilized powder			
RUNX	RUNX2			
AML3	AML3; CBFA1; OSF2; PEBP2A; Runt-related transcription factor 2; Acute myeloid			
leuker	mia 3 protein; Core-b	inding factor subunit alpha-1; CBF-alpha-1; Oncogene		
AML-3	3; Osteoblast-specific	transcription factor 2; OSF-2; Polyomavirus		
enhan	cer-binding protein	2		
860 (H	860 (Human)			
Q1395	Q13950 (Human)			
> 97%				
Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
appro	priate coupling effici	ency. The oligo is subsequently purified by affinity-solid		
phase	extraction. The ann	ealed RNA duplex is further analyzed by mass		
spectr	ometry to verify the	exact composition of the duplex. Each lot is compared to		
the pr	evious lot by mass s	pectrometry to ensure maximum lot-to-lot consistency.		
We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
huma	n RUNX2 gene. Each	vial contains 5 nmol of lyophilized siRNA. The duplexes		
can be	e transfected individu	ually or pooled together to achieve knockdown of the		
target	gene, which is most	commonly assessed by qPCR or western blot.		
	thetic siRNA RUNX knock Lyoph RUNX AML3 leuker AML-3 enhan 860 (H Q1399 > 97% Oligor appro phase spectr the pr We of huma can be	thetic H siRNA to inhibit RUNX2 exp RUNX2 siRNA (Human) is a knock down gene expressi Lyophilized powder RUNX2 AML3; CBFA1; OSF2; PEBP leukemia 3 protein; Core-b AML-3; Osteoblast-specific enhancer-binding protein 3 860 (Human) Q13950 (Human) > 97% Oligonucleotide synthesis a appropriate coupling effici phase extraction. The anne spectrometry to verify the the previous lot by mass sp We offers pre-designed set human RUNX2 gene. Each can be transfected individu		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

Component	15 nmol	30 nmol
RUNX2 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
RUNX2 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
RUNX2 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 µM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 µl
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 µl
		50 nM	5 μl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com