

# **Product Data Sheet**

### TMED2 siRNA (Human)

| Catalog #   | Source    | Reactivity  | Applicat                          | ions                            |  |  |
|---|-----------|---|-----------------------------------|---------------------------------|--|--|
| CRH7480   | Synthetic | н   | RNAi                              |                                 |  |  |
| Description   | siRNA     | to inhibit TMED2 ex   | pression using RNA interfe        | erence                          |  |  |
| Specificity   | TMED      | TMED2 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to    |                                   |                                 |  |  |
|   | knock     | down gene expressi  | on.                               |                                 |  |  |
| Form  | Lyoph     | Lyophilized powder  |                                   |                                 |  |  |
| Gene Symbol   | TMED      | TMED2   |                                   |                                 |  |  |
| Alternative N   | ames RNP2 | RNP24; Transmembrane emp24 domain-containing protein 2; Membrane protein              |                                   |                                 |  |  |
|   | p24A;     | ; p24; p24 family pro   | ein beta-1; p24beta1              |                                 |  |  |
| Entrez Gene   | 10959     | 10959 (Human)   |                                   |                                 |  |  |
| SwissProt   | Q153      | Q15363 (Human)  |                                   |                                 |  |  |
| Purity  | > 97%     | > 97%   |                                   |                                 |  |  |
| Quality Control Oligonucleotide synthesis is monitored base by base through tr                  |           |   | through trityl analysis to ensure |                                 |  |  |
|   | appro     | appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid |                                   |                                 |  |  |
| phase extraction. The annealed RNA duplex is further analyzed by ma                             |           |   | r analyzed by mass                |                                 |  |  |
|   | spect     | rometry to verify the   | exact composition of the          | duplex. Each lot is compared to |  |  |
|   | the pi    | revious lot by mass s   | pectrometry to ensure ma          | ximum lot-to-lot consistency.   |  |  |
| <b>Components</b> We offers pre-designed sets of 3 different target-specific siRNA oligo duplex |           |   | cific siRNA oligo duplexes of     |                                 |  |  |
|   | huma      | n TMED2 gene. Each  | vial contains 5 nmol of lyc       | philized siRNA. The duplexes    |  |  |
|   | can b     | e transfected individ   | ually or pooled together to       | achieve knockdown of the        |  |  |
|   | target    | target gene, which is most commonly assessed by qPCR or western blot.                 |                                   |                                 |  |  |
|   | Com       | ponent  | 15 nmol                           | 30 nmol                         |  |  |
|   | TME       | D2 siRNA (Human) -  | A 5 nmol x 1                      | . 5 nmol x 2                    |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



# **Product Data Sheet**

| TMED2 siRNA (Human) - B | 5 nmol x 1   | 5 nmol x 2   |
|-------------------------|--------------|--------------|
| TMED2 siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control        | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water              | 1 ml x 1     | 1 ml x 2     |

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 µl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 μl         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|         |              | 10 nM               | 0.25 μl       | 1 μl         |
|         |              | 100 nM              | 5 μl          | 2 μl         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μl         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
|         |              | 100 nM              | 10 µl         | 5 µl         |
| 6-well  | 2 ml         | 50 nM               | 5 µl          | 5 μΙ         |
|         |              | 10 nM               | 1 μΙ          | 5 μΙ         |

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |