

Product Data Sheet

KIF2C siRNA (Human)

| Catalog # Sour | ce Reactivity | Applications | | |
|-------------------|---|---------------------------------------|-----------------------|--|
| CRH7514 Synt | hetic H | RNAi | | |
| Description | siRNA to inhibit KIF2C expre | ssion using RNA interference | | |
| Specificity | KIF2C siRNA (Human) is a tai | rget-specific 19-23 nt siRNA oligo d | luplexes designed to | |
| | knock down gene expressior | ۱. | | |
| Form | Lyophilized powder | | | |
| Gene Symbol | KIF2C | | | |
| Alternative Names | KNSL6; Kinesin-like protein k | (IF2C; Kinesin-like protein 6; Mitoti | ic | |
| | centromere-associated kines | sin; MCAK | | |
| Entrez Gene | 11004 (Human) | | | |
| SwissProt | Q99661 (Human) | | | |
| Purity | > 97% | | | |
| Quality Control | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appropriate coupling efficier | ncy. The oligo is subsequently purif | ied by affinity-solid | |
| | phase extraction. The annealed RNA duplex is further analyzed by mass | | | |
| | spectrometry to verify the e | xact composition of the duplex. Ea | ch lot is compared to | |
| | the previous lot by mass spe | ectrometry to ensure maximum lot | -to-lot consistency. | |
| Components | Components We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes | | | |
| | human KIF2C gene. Each via | l contains 5 nmol of lyophilized siR | NA. The duplexes can | |
| | be transfected individually o | r pooled together to achieve knoc | kdown of the target | |
| | gene, which is most commonly assessed by qPCR or western blot. | | | |
| Component 15 nmol | | | 30 nmol | |
| | KIF2C siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| KIF2C siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
|-------------------------|--------------|--------------|
| KIF2C siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 µl |
| | | 10 nM | 0.25 μl | 1 µl |
| | | 100 nM | 5 μl | 2 μl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μl |
| | | 100 nM | 10 µl | 5 μΙ |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μΙ |
| | | 10 nM | 1 μl | 5 μΙ |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |