

## HHLA3 siRNA (Human)

Catalog #	Source	Reactivity	Applications
CRH7629	Synthetic	H	RNAi
<b>Description</b>	siRNA to inhibit HHLA3 expression using RNA interference		
<b>Specificity</b>	HHLA3 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to knock down gene expression.		
<b>Form</b>	Lyophilized powder		
<b>Gene Symbol</b>	HHLA3		
<b>Alternative Names</b>	HERV-H LTR-associating protein 3; Human endogenous retrovirus-H long terminal repeat-associating protein 3		
<b>Entrez Gene</b>	11147 (Human)		
<b>SwissProt</b>	Q9XRX5 (Human)		
<b>Purity</b>	> 97%		
<b>Quality Control</b>	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid phase extraction. The annealed RNA duplex is further analyzed by mass spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.		
<b>Components</b>	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of human HHLA3 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can be transfected individually or pooled together to achieve knockdown of the target gene, which is most commonly assessed by qPCR or western blot.		

Component	15 nmol	30 nmol
HHLA3 siRNA (Human) - A	5 nmol x 1	5 nmol x 2

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

### COHESION BIOSCIENCES LIMITED

**WEB**  
www.cohesionbio.com

**ORDER**  
order@cohesionbio.com

**SUPPORT**  
techsupport@cohesionbio.com

**CUSTOM**  
custom@cohesionbio.com

## Product Data Sheet

HHLA3 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
HHLA3 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

### Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume of medium	Final concentration of siRNA	siRNA (20 $\mu$ M)	Lipofectamin 2000
96-well	100 $\mu$ l	100 nM	0.5 $\mu$ l	0.25 $\mu$ l
		50 nM	0.25 $\mu$ l	0.25 $\mu$ l
		10 nM	0.05 $\mu$ l	0.25 $\mu$ l
24-well	500 $\mu$ l	100 nM	2.5 $\mu$ l	1 $\mu$ l
		50 nM	1.25 $\mu$ l	1 $\mu$ l
		10 nM	0.25 $\mu$ l	1 $\mu$ l
12-well	1 ml	100 nM	5 $\mu$ l	2 $\mu$ l
		50 nM	2.5 $\mu$ l	2 $\mu$ l
		10 nM	0.5 $\mu$ l	2 $\mu$ l
6-well	2 ml	100 nM	10 $\mu$ l	5 $\mu$ l
		50 nM	5 $\mu$ l	5 $\mu$ l
		10 nM	1 $\mu$ l	5 $\mu$ l

### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

**Application key:** E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC- Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference

**Species reactivity key:** H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb- Rabbit, S- Sheep, Z- Zebrafish

## COHESION BIOSCIENCES LIMITED

### WEB

[www.cohesionbio.com](http://www.cohesionbio.com)

### ORDER

[order@cohesionbio.com](mailto:order@cohesionbio.com)

### SUPPORT

[techsupport@cohesionbio.com](mailto:techsupport@cohesionbio.com)

### CUSTOM

[custom@cohesionbio.com](mailto:custom@cohesionbio.com)