

# **Product Data Sheet**

## **CELF3 siRNA (Human)**

| Catalog #      | Source    | Reactivity   | Applications  |  |  |
|----------------|-----------|--|---|--|--|
| CRH7669        | Synthetic | н  | RNAi  |  |  |
| Description    | siRNA     | to inhibit CELF3 exp   | ression using RNA interference                              |  |  |
| Specificity    | CELFS     | 8 siRNA (Human) is a   | target-specific 19-23 nt siRNA oligo duplexes designed to   |  |  |
|                | knock     | down gene expressi   | on.   |  |  |
| Form           | Lyoph     | nilized powder   |   |  |  |
| Gene Symbol    | CELFS     | 3  |   |  |  |
| Alternative Na | ames BRUN | BRUNOL1; CAGH4; ERDA4; TNRC4; CUGBP Elav-like family member 3; CELF-3; |   |  |  |
|                | Brund     | o-like protein 1; CAG  | repeat protein 4; CUG-BP- and ETR-3-like factor 3;          |  |  |
|                | ELAV-     | type RNA-binding pro   | otein 1; ETR-1; Expanded repeat domain protein CAG/CTG      |  |  |
|                | 4; RN     | A-binding protein BR   | UNOL-1; Trinucleotide repeat-containing gene 4 protein      |  |  |
| Entrez Gene    | 1118      | 9 (Human)  |   |  |  |
| SwissProt      | Q5SZ      | Q8 (Human)   |   |  |  |
| Purity         | > 97%     | 0  |   |  |  |
| Quality Contro | ol Oligo  | nucleotide synthesis   | is monitored base by base through trityl analysis to ensure |  |  |
|                | appro     | priate coupling effici   | ency. The oligo is subsequently purified by affinity-solid  |  |  |
|                | phase     | e extraction. The ann  | ealed RNA duplex is further analyzed by mass                |  |  |
|                | spect     | rometry to verify the  | exact composition of the duplex. Each lot is compared to    |  |  |
|                | the p     | revious lot by mass s  | pectrometry to ensure maximum lot-to-lot consistency.       |  |  |
| Components     | We o      | ffers pre-designed se  | ts of 3 different target-specific siRNA oligo duplexes of   |  |  |
|                | huma      | in CELF3 gene. Each v  | ial contains 5 nmol of lyophilized siRNA. The duplexes can  |  |  |
|                | be tra    | ansfected individually   | or pooled together to achieve knockdown of the target       |  |  |
|                | gene,     | which is most comm   | only assessed by qPCR or western blot.                      |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

| Component               | 15 nmol      | 30 nmol      |
|-------------------------|--------------|--------------|
| CELF3 siRNA (Human) - A | 5 nmol x 1   | 5 nmol x 2   |
| CELF3 siRNA (Human) - B | 5 nmol x 1   | 5 nmol x 2   |
| CELF3 siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control        | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water              | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 μl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 μΙ         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|         |              | 10 nM               | 0.25 μl       | 1 μΙ         |
|         |              | 100 nM              | 5 µl          | 2 μΙ         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
| 6-well  | 2 ml         | 100 nM              | 10 µl         | 5 µl         |
|         |              | 50 nM               | 5 µl          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

 WEB
 ORDER
 SUPPORT
 CUSTOM

 www.cohesionbio.com
 order@cohesionbio.com
 techsupport@cohesionbio.com
 custom@cohesionbio.com