

# **Product Data Sheet**

## **INPP5F siRNA (Human)**

| Catalog #       | Source     | Reactivity  | Applications  |  |  |  |
|-----------------|------------|---|---|--|--|--|
| CRH7833         | Synthetic  | н   | RNAi  |  |  |  |
| Description     | siRNA      | to inhibit INPP5F ex  | pression using RNA interference                             |  |  |  |
| Specificity     | INPP5      | INPP5F siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to |   |  |  |  |
|                 | knock      | k down gene expressi  | on.   |  |  |  |
| Form            | Lyoph      | nilized powder  |   |  |  |  |
| Gene Symbol     | INPP5      | INPP5F  |   |  |  |  |
| Alternative Na  | ames KIAA( | KIAA0966; SAC2; Phosphatidylinositide phosphatase SAC2; Inositol polyphosphate      |   |  |  |  |
|                 | 5-phc      | osphatase F; Sac dom  | ain-containing inositol phosphatase 2; Sac                  |  |  |  |
|                 | doma       | in-containing phosph  | noinositide 5-phosphatase 2; hSAC2                          |  |  |  |
| Entrez Gene     | 22876      | 6 (Human)   |   |  |  |  |
| SwissProt       | Q9Y2       | Q9Y2H2 (Human)  |   |  |  |  |
| Purity          | > 97%      | > 97%   |   |  |  |  |
| Quality Control | ol Oligo   | nucleotide synthesis  | is monitored base by base through trityl analysis to ensure |  |  |  |
|                 | appro      | opriate coupling effici   | ency. The oligo is subsequently purified by affinity-solid  |  |  |  |
|                 | phase      | e extraction. The ann   | ealed RNA duplex is further analyzed by mass                |  |  |  |
|                 | spect      | rometry to verify the   | exact composition of the duplex. Each lot is compared to    |  |  |  |
|                 | the p      | revious lot by mass s   | pectrometry to ensure maximum lot-to-lot consistency.       |  |  |  |
| Components      | We of      | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of  |   |  |  |  |
|                 | huma       | human INPP5F gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes     |   |  |  |  |
|                 | can b      | e transfected individu  | ually or pooled together to achieve knockdown of the        |  |  |  |
|                 | target     | target gene, which is most commonly assessed by qPCR or western blot.               |   |  |  |  |
|                 | Com        | iponent   | 15 nmol 30 nmol   |  |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



# **Product Data Sheet**

| INPP5F siRNA (Human) - A | 5 nmol x 1   | 5 nmol x 2   |
|--------------------------|--------------|--------------|
| INPP5F siRNA (Human) - B | 5 nmol x 1   | 5 nmol x 2   |
| INPP5F siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control         | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water               | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 µl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 μl         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|         |              | 10 nM               | 0.25 μl       | 1 µl         |
|         |              | 100 nM              | 5 μl          | 2 μl         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
|         |              | 100 nM              | 10 µl         | 5 µl         |
| 6-well  | 2 ml         | 50 nM               | 5 µl          | 5 µl         |
|         |              | 10 nM               | 1 μl          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

## **Product Data Sheet**

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com