

# **Product Data Sheet**

### SETX siRNA (Human)

| Catalog #     | Source     | Reactivity  | Applications                      |                          |  |
|---------------|------------|---|-----------------------------------|--------------------------|--|
| CRH7971       | Synthetic  | н   | RNAi                              |                          |  |
| Description   | siRNA      | to inhibit SETX expres  | ssion using RNA interference      |                          |  |
| Specificity   | SETX s     | SETX siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to     |                                   |                          |  |
|               | knock      | down gene expressio   | n.                                |                          |  |
| Form          | Lyoph      | ilized powder   |                                   |                          |  |
| Gene Symbol   | SETX       | SETX  |                                   |                          |  |
| Alternative N | ames ALS4; | ALS4; KIAA0625; SCAR1; Probable helicase senataxin; Amyotrophic lateral sclerosis 4   |                                   |                          |  |
|               | protei     | n; SEN1 homolog   |                                   |                          |  |
| Entrez Gene   | 23064      | (Human)   |                                   |                          |  |
| SwissProt     | Q7Z33      | Q7Z333 (Human)  |                                   |                          |  |
| Purity        | > 97%      | > 97%   |                                   |                          |  |
| Quality Contr | ol Oligor  | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure |                                   |                          |  |
|               | appro      | priate coupling efficie   | ncy. The oligo is subsequently pu | rified by affinity-solid |  |
|               | phase      | phase extraction. The annealed RNA duplex is further analyzed by mass                 |                                   |                          |  |
|               | spectr     | spectrometry to verify the exact composition of the duplex. Each lot is compared to   |                                   |                          |  |
|               | the pr     | the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.       |                                   |                          |  |
| Components    | We of      | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of    |                                   |                          |  |
|               | huma       | human SETX gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can     |                                   |                          |  |
|               | be tra     | be transfected individually or pooled together to achieve knockdown of the target     |                                   |                          |  |
|               | gene,      | gene, which is most commonly assessed by qPCR or western blot.                        |                                   |                          |  |
|               | Com        | ponent  | 15 nmol                           | 30 nmol                  |  |
|               | SETX       | siRNA (Human) - A   | 5 nmol x 1                        | 5 nmol x 2               |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



## **Product Data Sheet**

| SETX siRNA (Human) - B | 5 nmol x 1   | 5 nmol x 2   |
|------------------------|--------------|--------------|
| SETX siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control       | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water             | 1 ml x 1     | 1 ml x 2     |

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 µl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 μl         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|         |              | 10 nM               | 0.25 μl       | 1 μΙ         |
|         |              | 100 nM              | 5 μl          | 2 μl         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
|         |              | 100 nM              | 10 µl         | 5 μΙ         |
| 6-well  | 2 ml         | 50 nM               | 5 µl          | 5 μΙ         |
|         |              | 10 nM               | 1 μΙ          | 5 μΙ         |

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |