

Product Data Sheet

TMED3 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | | |
|--|------------|---|-----------------------------------|------------------------|--|--|
| CRH8260 | Synthetic | н | RNAi | | | |
| Description | siRNA | to inhibit TMED3 ex | pression using RNA interference | | | |
| Specificity | TMED | TMED3 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | | |
| | knock | knock down gene expression. | | | | |
| Form | Lyoph | Lyophilized powder | | | | |
| Gene Symbol | TMED | TMED3 | | | | |
| Alternative N | ames C15or | C15orf22; Transmembrane emp24 domain-containing protein 3; Membrane protein | | | | |
| | p24B; | p24 family protein g | amma-4; p24gamma4; p26 | | | |
| Entrez Gene | 23423 | 23423 (Human) | | | | |
| SwissProt | Q9Y3 | Q9Y3Q3 (Human) | | | | |
| Purity | > 97% | > 97% | | | | |
| Quality Control Oligonucleotide synthesis is monitored base by base through trityl analys | | | igh trityl analysis to ensure | | | |
| | appro | appropriate coupling efficiency. The oligo is subsequently purified by affinity-so | | | | |
| | phase | phase extraction. The annealed RNA duplex is further analyzed by mass | | | | |
| | spect | spectrometry to verify the exact composition of the duplex. Each lot is compared to | | | | |
| | the pr | the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency. | | | | |
| Components We offers pre-designed sets of 3 different target-specific siRNA oligo duple | | | iRNA oligo duplexes of | | | |
| | huma | n TMED3 gene. Each | vial contains 5 nmol of lyophiliz | ed siRNA. The duplexes | | |
| | can be | e transfected individ | ually or pooled together to achie | eve knockdown of the | | |
| | target | target gene, which is most commonly assessed by qPCR or western blot. | | | | |
| | Com | ponent | 15 nmol | 30 nmol | | |
| | TME | D3 siRNA (Human) - | A 5 nmol x 1 | 5 nmol x 2 | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| TMED3 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
|-------------------------|--------------|--------------|
| TMED3 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| _ | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| _ | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 μl | 2 μl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| _ | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μΙ |
| | | 10 nM | 1 µl | 5 μΙ |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |