

# **Product Data Sheet**

## PSD4 siRNA (Human)

ic H	RNAi	
siRNA to inhibit PSD4 ex	pression using RNA interference	
PSD4 siRNA (Human) is a	a target-specific 19-23 nt siRNA oligo duplexes designed to	
knock down gene expres	ssion.	
Lyophilized powder		
Gene Symbol PSD4		
EFA6B; TIC; PH and SEC7	domain-containing protein 4; Exchange factor for	
ADP-ribosylation factor §	guanine nucleotide factor 6 B; Exchange factor for ARF6 B;	
Pleckstrin homology and	SEC7 domain-containing protein 4; Telomeric of	
interleukin-1 cluster pro	tein	
23550 (Human)		
Q8NDX1 (Human)		
> 97%		
Oligonucleotide synthes	is is monitored base by base through trityl analysis to ensure	
appropriate coupling eff	iciency. The oligo is subsequently purified by affinity-solid	
phase extraction. The ar	nealed RNA duplex is further analyzed by mass	
spectrometry to verify th	ne exact composition of the duplex. Each lot is compared to	
the previous lot by mass	spectrometry to ensure maximum lot-to-lot consistency.	
We offers pre-designed	sets of 3 different target-specific siRNA oligo duplexes of	
human PSD4 gene. Each	vial contains 5 nmol of lyophilized siRNA. The duplexes can	
be transfected individua	lly or pooled together to achieve knockdown of the target	
gene, which is most com	monly assessed by qPCR or western blot.	
	siRNA to inhibit PSD4 ex PSD4 siRNA (Human) is a knock down gene express Lyophilized powder PSD4 EFA6B; TIC; PH and SEC7 ADP-ribosylation factor g Pleckstrin homology and interleukin-1 cluster pro- 23550 (Human) Q8NDX1 (Human) > 97% Oligonucleotide synthes appropriate coupling eff phase extraction. The ar spectrometry to verify th the previous lot by mass We offers pre-designed s human PSD4 gene. Each be transfected individua	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

Component	15 nmol	30 nmol
PSD4 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
PSD4 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
PSD4 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
96-well	100 μl	100 nM	0.5 μl	0.25 μl
		50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
24-well	500 μl	100 nM	2.5 μl	1 μl
		50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
12-well 1		100 nM	5 µl	2 μl
	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 µl
	2 ml	50 nM	5 µl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com