

# **Product Data Sheet**

## MGAT4C siRNA (Human)

Catalog #	Source	Reactivity	Applications
CRH8531	Synthetic	Н	RNAi
Description	siRNA	to inhibit MGAT4C e	pression using RNA interference
Specificity	MGA	T4C siRNA (Human) is	a target-specific 19-23 nt siRNA oligo duplexes designed
	to kn	ock down gene expre	ssion.
Form	Lyopł	nilized powder	
Gene Symbol	MGA	T4C	
Alternative Na	ames Alpha	a-1.3-mannosyl-glycop	protein 4-beta-N-acetylglucosaminyltransferase C;
	N-ace	etylglucosaminyltrans	erase IV homolog; hGnT-IV-H;
	N-gly	cosyl-oligosaccharide	glycoprotein N-acetylglucosaminyltransferase IVc;
	GlcNA	Ac-T IVc; GnT-IVc; N-ad	etylglucosaminyltransferase IVc;
	UDP-	N-acetylglucosamine:	alpha-1.3-D-mannoside
	beta-	1.4-N-acetylglucosam	inyltransferase IVc
Entrez Gene	25834	4 (Human)	
SwissProt	Q9UE	3M8 (Human)	
Purity	> 97%	6	
Quality Contro	ol Oligo	nucleotide synthesis i	s monitored base by base through trityl analysis to ensure
	appro	opriate coupling efficie	ency. The oligo is subsequently purified by affinity-solid
	phase	e extraction. The anne	aled RNA duplex is further analyzed by mass
	spect	rometry to verify the	exact composition of the duplex. Each lot is compared to
	the p	revious lot by mass sp	ectrometry to ensure maximum lot-to-lot consistency.
Components	We o	ffers pre-designed set	s of 3 different target-specific siRNA oligo duplexes of
	huma	an MGAT4C gene. Eacl	n vial contains 5 nmol of lyophilized siRNA. The duplexes

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

can be transfected individually or pooled together to achieve knockdown of the

target gene, which is most commonly assessed by qPCR or western blot.

Component	15 nmol	30 nmol
MGAT4C siRNA (Human) - A	5 nmol x 1	5 nmol x 2
MGAT4C siRNA (Human) - B	5 nmol x 1	5 nmol x 2
MGAT4C siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 µl of DEPC water to get a final concentration of 20 µM.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



For research purposes only, not for human use

# **Product Data Sheet**

		100 nM	10 µl	5 μl
6-well	2 ml	50 nM	5 µl	5 µl
		10 nM	1 µl	5 µl

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com