

Product Data Sheet

GIT1 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | |
|---------------|------------|--|--------------------------------------|---------------------------|--|
| CRH9131 | Synthetic | н | RNAi | | |
| Description | siRNA | A to inhibit GIT1 expre | ssion using RNA interference | | |
| Specificity | GIT1 | siRNA (Human) is a ta | rget-specific 19-23 nt siRNA oligo | duplexes designed to | |
| | knock | < down gene expressio | on. | | |
| Form | Lyoph | nilized powder | | | |
| Gene Symbol | GIT1 | | | | |
| Alternative N | ames ARF C | ARF GTPase-activating protein GIT1; ARF GAP GIT1; Cool-associated and | | | |
| | tyrosi | ine-phosphorylated pr | otein 1; CAT-1; CAT1; G protein-cc | oupled receptor | |
| | kinas | e-interactor 1; GRK-in | teracting protein 1 | | |
| Entrez Gene | 28964 | 4 (Human) | | | |
| SwissProt | Q9Y2 | Q9Y2X7 (Human) | | | |
| Purity | > 97% | 6 | | | |
| Quality Contr | ol Oligo | nucleotide synthesis is | s monitored base by base through | trityl analysis to ensure | |
| | appro | opriate coupling efficie | ency. The oligo is subsequently pu | rified by affinity-solid | |
| | phase | e extraction. The anne | aled RNA duplex is further analyze | ed by mass | |
| | spect | rometry to verify the | exact composition of the duplex. I | Each lot is compared to | |
| | the p | revious lot by mass sp | ectrometry to ensure maximum lo | ot-to-lot consistency. | |
| Components | We o | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | huma | an GIT1 gene. Each via | l contains 5 nmol of lyophilized sif | RNA. The duplexes can | |
| | be tra | ansfected individually | or pooled together to achieve kno | ockdown of the target | |
| | gene, | which is most commo | only assessed by qPCR or western | blot. | |
| | Com | ponent | 15 nmol | 30 nmol | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| GIT1 siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
|------------------------|--------------|--------------|
| GIT1 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| GIT1 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μl |
| | | 100 nM | 5 μl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μl |
| | | 10 nM | 1 μl | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com