

Product Data Sheet

GMPPA siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | |
|---|-----------|---|-----------------------------------|----------------------------|--|
| CRH9273 | Synthetic | Н | RNAi | | |
| Description | siRNA | to inhibit GMPPA ex | pression using RNA interference | 2 | |
| Specificity | GMPF | GMPPA siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | |
| | knock | down gene expressi | on. | | |
| Form | Lyoph | ilized powder | | | |
| Gene Symbol | GMPF | GMPPA | | | |
| Alternative N | ames Mann | Mannose-1-phosphate guanyltransferase alpha; GDP-mannose pyrophosphorylase | | | |
| | A; GN | 1PP-alpha; GTP-manr | nose-1-phosphate guanylyltransf | ferase alpha | |
| Entrez Gene | 29926 | 29926 (Human) | | | |
| SwissProt | Q96IJ | Q96IJ6 (Human) | | | |
| Purity > 97% | | | | | |
| Quality Control Oligonucleotide synthesis is monitored base by base through trityl analys | | | igh trityl analysis to ensure | | |
| | appro | priate coupling effici | ency. The oligo is subsequently (| purified by affinity-solid | |
| | phase | phase extraction. The annealed RNA duplex is further analyzed by mass | | | |
| | spect | spectrometry to verify the exact composition of the duplex. Each lot is compared to | | | |
| | the pr | the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency. | | | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | huma | human GMPPA gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes | | | |
| | can be | can be transfected individually or pooled together to achieve knockdown of the | | | |
| | target | target gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol | 30 nmol | |
| | GMP | PPA siRNA (Human) - | A 5 nmol x 1 | 5 nmol x 2 | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| GMPPA siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
|-------------------------|--------------|--------------|
| GMPPA siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μl |
| | | 100 nM | 5 μl | 2 µl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μΙ |
| | | 10 nM | 1 μl | 5 μΙ |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |