

# **Product Data Sheet**

### ZNF639 siRNA (Human)

Catalog #	Source	Reactivity	Application	S		
CRH9596	Synthetic	н	RNAi			
Description	siRNA	to inhibit ZNF639 ex	pression using RNA interferer	nce		
Specificity	ZNF63	ZNF639 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene expression	on.			
Form	Lyoph	ilized powder				
Gene Symbol	ZNF63	ZNF639				
Alternative N	ames ZASC1	ZASC1; Zinc finger protein 639; Zinc finger protein ANC_2H01; Zinc finger protein				
	ZASC1	L				
Entrez Gene	51193	51193 (Human)				
SwissProt	Q9UII	Q9UID6 (Human)				
Purity	> 97%					
Quality Contr	Unality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to			ough trityl analysis to ensure		
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-sol				
phase extraction. The annealed RNA duplex is further analyzed by mass			nalyzed by mass			
	specti	rometry to verify the	exact composition of the dup	blex. Each lot is compared to		
	the pr	revious lot by mass sp	pectrometry to ensure maxim	um lot-to-lot consistency.		
<b>Components</b> We offers pre-designed sets of 3 different target-specific siRNA oligo duple			c siRNA oligo duplexes of			
	huma	n ZNF639 gene. Each	vial contains 5 nmol of lyoph	ilized siRNA. The duplexes		
	can be	e transfected individu	ally or pooled together to ac	hieve knockdown of the		
	target	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	ZNF6	539 siRNA (Human) -	A 5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



# **Product Data Sheet**

ZNF639 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
ZNF639 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
_		10 nM	0.25 μl	1 μΙ
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 µl	5 µl
		10 nM	1 μΙ	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com