

Product Data Sheet

TRPV2 siRNA (Human)

Catalog #	Source	Reactivity	Applications			
CRH9744	Synthetic	н	RNAi			
Description	siRNA	to inhibit TRPV2 exp	pression using RNA interference			
Specificity	TRPV2	TRPV2 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene expressi	on.			
Form	Lyoph	Lyophilized powder				
Gene Symbol	TRPV2	TRPV2				
Alternative N	ames VRL; 1	VRL; Transient receptor potential cation channel subfamily V member 2; TrpV2;				
	Osm-9	9-like TRP channel 2;	OTRPC2; Vanilloid receptor-like	protein 1; VRL-1		
Entrez Gene	51393	51393 (Human)				
SwissProt	Q9Y5	Q9Y5S1 (Human)				
Purity	> 97%	> 97%				
Quality Control Oligonucleotide synthesis is monitored base by base through trityl ana			gh trityl analysis to ensure			
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-so				
	phase	e extraction. The ann	ealed RNA duplex is further analy	vzed by mass		
	spect	rometry to verify the	exact composition of the duplex	. Each lot is compared to		
	the pi	revious lot by mass s	pectrometry to ensure maximum	lot-to-lot consistency.		
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	huma	n TRPV2 gene. Each	vial contains 5 nmol of lyophilize	d siRNA. The duplexes can		
	be tra	be transfected individually or pooled together to achieve knockdown of the target				
	gene,	gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	TRP	/2 siRNA (Human) - A	5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

TRPV2 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
TRPV2 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 µl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com