

# **Product Data Sheet**

## MED15 siRNA (Human)

Catalog #	Source	Reactivity	Applications			
CRH9845	Synthetic	н	RNAi			
Description	siRN	A to inhibit MED15 ex	pression using RNA interference			
Specificity	MED	15 siRNA (Human) is	a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knoc	k down gene expressi	on.			
Form	Lyop	hilized powder				
Gene Symbol	MED	15				
Alternative Na	ames ARC1	.05; CTG7A; PCQAP; T	IG1; TNRC7; Mediator of RNA polymerase II transcription			
	subu	nit 15; Activator-recru	ited cofactor 105 kDa component; ARC105; CTG repeat			
	prote	ein 7a; Mediator com	blex subunit 15; Positive cofactor 2			
	gluta	mine/Q-rich-associat	ed protein; PC2 glutamine/Q-rich-associated protein;			
	TPA-i	nducible gene 1 prote	ein; TIG-1; Trinucleotide repeat-containing gene 7 protein			
Entrez Gene	5158	6 (Human)				
SwissProt	Q96F	Q96RN5 (Human)				
Purity	> 97%	> 97%				
Quality Control	ol Oligo	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
	appro	opriate coupling effici	ency. The oligo is subsequently purified by affinity-solid			
	phas	e extraction. The ann	ealed RNA duplex is further analyzed by mass			
	spect	trometry to verify the	exact composition of the duplex. Each lot is compared to			
	the p	revious lot by mass s	pectrometry to ensure maximum lot-to-lot consistency.			
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	huma	human MED15 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes				
	can b	e transfected individ	ually or pooled together to achieve knockdown of the			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

target gene, which is most commonly assessed by qPCR or western blot.	
---	--

Component	15 nmol	30 nmol
MED15 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
MED15 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
MED15 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
	100 µl	100 nM	0.5 μl	0.25 μl
96-well		50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
24-well	500 μl	100 nM	2.5 μl	1 µl
		50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
12-well	1 ml	100 nM	5 μl	2 μΙ
		50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



For research purposes only, not for human use

# **Product Data Sheet**

50 nM5 μl5 μl10 nM1 μl5 μl	o. /o. 1.1				
50 nM 5 μl 5 μl		10	) nM	1 µl	5 μl
		50	) nM	5 μl	5 μΙ

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com