

Product Data Sheet

TMEM66 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | |
|---------------|-----------|---|--|--|--|
| CRH9894 | Synthetic | н | RNAi | | |
| Description | siRNA | to inhibit TMEM66 | expression using RNA interference | | |
| Specificity | TMEN | /166 siRNA (Human) i | s a target-specific 19-23 nt siRNA oligo duplexes designed | | |
| | to kno | ock down gene expre | ssion. | | |
| Form | Lyoph | ilized powder | | | |
| Gene Symbol | TMEN | TMEM66 | | | |
| Alternative N | ames SARA | SARAF; XTP3; Store-operated calcium entry-associated regulatory factor; SARAF; | | | |
| | SOCE | -associated regulator | y factor; HBV X-transactivated gene 3 protein; HBV | | |
| | XAg-t | ransactivated proteir | a 3; Protein FOAP-7; Transmembrane protein 66 | | |
| Entrez Gene | 51669 |) (Human) | | | |
| SwissProt | Q96B | Q96BY9 (Human) | | | |
| Purity | > 97% | ,) | | | |
| Quality Contr | ol Oligoi | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure | | | |
| | appro | priate coupling effici | ency. The oligo is subsequently purified by affinity-solid | | |
| | phase | extraction. The ann | ealed RNA duplex is further analyzed by mass | | |
| | spect | rometry to verify the | exact composition of the duplex. Each lot is compared to | | |
| | the pi | revious lot by mass s | pectrometry to ensure maximum lot-to-lot consistency. | | |
| Components | We of | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of | | | |
| | huma | n TMEM66 gene. Eac | ch vial contains 5 nmol of lyophilized siRNA. The duplexes | | |
| | can b | e transfected individu | ually or pooled together to achieve knockdown of the | | |
| | target | target gene, which is most commonly assessed by qPCR or western blot. | | | |
| | Com | ponent | 15 nmol 30 nmol | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | СИЗТОМ |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| TMEM66 siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
|--------------------------|--------------|--------------|
| TMEM66 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| TMEM66 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μΙ |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 µl | 2 μΙ |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μl |
| | | 10 nM | 1 μΙ | 5 μΙ |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com