

Product Data Sheet

FAM63B siRNA (Human)

Catalog #	Source	Reactivity	Арр	ications	
CRJ0163	Synthetic	H	RNA		
Description	siRNA	to inhibit FAM63B e	xpression using RNA ir	iterference	
Specificity	FAM6	3B siRNA (Human) is	a target-specific 19-23	3 nt siRNA oligo duplexes des	igned
	to kno	ock down gene expre	ssion.		
Form	Lyoph	ilized powder			
Gene Symbol	FAM6	3B			
Alternative N	ames KIAA1	164; Protein FAM63I	3		
Entrez Gene	54629) (Human)			
SwissProt	Q8NB	R6 (Human)			
Purity	> 97%	,)			
Quality Conti	ol Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	priate coupling effici	ency. The oligo is subs	equently purified by affinity-	solid
	phase	e extraction. The anno	ealed RNA duplex is fu	rther analyzed by mass	
	spect	rometry to verify the	exact composition of	the duplex. Each lot is compa	red to
	the p	revious lot by mass s	pectrometry to ensure	maximum lot-to-lot consiste	ncy.
Components	We of	ffers pre-designed se	ts of 3 different target	specific siRNA oligo duplexes	s of
	huma	n FAM63B gene. Eacl	n vial contains 5 nmol	of lyophilized siRNA. The dup	lexes
	can b	e transfected individu	ally or pooled togeth	er to achieve knockdown of t	he
	target gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nm	ol 30 nmol	
	FAM	63B siRNA (Human) -	A 5 nmc	l x 1 5 nmol x 2	
	FAM	63B siRNA (Human) -	B 5 nmc	l x 1 5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

FAM	63B siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Nega	ative Control	2.5 nmol x 1	2.5 nmol x 2
DEP	C Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com