

# **Product Data Sheet**

### ST6GALNAC1 siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRJ0938	Synthetic	н	RNAi		
Description	siRNA	A to inhibit ST6GALN	AC1 expression using RNA interference	2	
Specificity	ST6G	ALNAC1 siRNA (Hum	an) is a target-specific 19-23 nt siRNA o	oligo duplexes	
	desig	ned to knock down g	ene expression.		
Form	Lyoph	nilized powder			
Gene Symbol	ST6G	ST6GALNAC1			
Alternative N	ames SIAT7	SIAT7A; Alpha-N-acetylgalactosaminide alpha-2.6-sialyltransferase 1; GalNAc			
	alpha	a-2.6-sialyltransferase	e I; ST6GalNAc I; ST6GalNAcI; Sialyltran	sferase 7A; SIAT7-A	
Entrez Gene	5580	8 (Human)			
SwissProt	Q9NS	SC7 (Human)			
Purity >		> 97%			
Quality Control Oligonucleotide synthesis is monitored base by base through trityl			tyl analysis to ensure		
	appro	opriate coupling effic	iency. The oligo is subsequently purifie	ed by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the p	the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.			
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	human ST6GALNAC1 gene. Each vial contains 5 nmol of lyophilized siRNA. The			
	duple	duplexes can be transfected individually or pooled together to achieve knockdown			
	of the	of the target gene, which is most commonly assessed by qPCR or western blot.			
	Com	nponent	15 nmol	30 nmol	
	ST60	GALNAC1 siRNA (Hur	nan) - A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com

## **Product Data Sheet**

DEPC Water	1 ml x 1	1 ml x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
ST6GALNAC1 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
ST6GALNAC1 siRNA (Human) - B	5 nmol x 1	5 nmol x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 µl	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com