

Product Data Sheet

TAOK1 siRNA (Human)

e	Reactivity	Applications		
etic	Н	RNAi		
siRNA to	inhibit TAOK1 ex	pression using RNA interference		
TAOK1 si	RNA (Human) is	a target-specific 19-23 nt siRNA oligo duplexes designed to		
knock do	own gene express	ion.		
Lyophiliz	ed powder			
TAOK1				
KIAA136	KIAA1361; MAP3K16; MARKK; Serine/threonine-protein kinase TAO1; Kinase from			
chicken ł	nomolog B; hKFC	-B; MARK Kinase; MARKK; Prostate-derived sterile 20-like		
kinase 2;	PSK-2; PSK2; Pro	ostate-derived STE20-like kinase 2; Thousand and one		
amino ao	cid protein kinase	e 1; TAOK1; hTAOK1		
57551 (H	luman)			
Q7L7X3 ((Human)			
> 97%				
Oligonuc	leotide synthesis	is monitored base by base through trityl analysis to ensure		
appropri	ate coupling effic	ciency. The oligo is subsequently purified by affinity-solid		
phase ex	traction. The anr	nealed RNA duplex is further analyzed by mass		
spectron	netry to verify th	e exact composition of the duplex. Each lot is compared to		
the previ	ious lot by mass s	spectrometry to ensure maximum lot-to-lot consistency.		
We offer	s pre-designed se	ets of 3 different target-specific siRNA oligo duplexes of		
human T	AOK1 gene. Each	vial contains 5 nmol of lyophilized siRNA. The duplexes		
can be tr	ansfected individ	lually or pooled together to achieve knockdown of the		
target ge	ene, which is mos	t commonly assessed by qPCR or western blot.		
	etic siRNA to TAOK1 si knock do Lyophiliz TAOK1 KIAA136 chicken l kinase 2; amino ad 57551 (H Q7L7X3 > 97% Oligonuc appropri phase ex spectron the prev We offer human T can be tr	etic H siRNA to inhibit TAOK1 ex TAOK1 siRNA (Human) is a knock down gene express Lyophilized powder TAOK1 KIAA1361; MAP3K16; MA chicken homolog B; hKFC kinase 2; PSK-2; PSK2; Pro amino acid protein kinase 57551 (Human) Q7L7X3 (Human) > 97% Oligonucleotide synthesis appropriate coupling effic phase extraction. The ann spectrometry to verify the the previous lot by mass s We offers pre-designed se human TAOK1 gene. Each can be transfected individ		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Component	15 nmol	30 nmol
TAOK1 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
TAOK1 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
TAOK1 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μl
		100 nM	5 µl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 µl
		50 nM	5 µl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com