

Product Data Sheet

DDX55 siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRJ1639	Synthetic	Н	RNAi		
Description	siRN	siRNA to inhibit DDX55 expression using RNA interference			
Specificity	DDX5	DDX55 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knoc	k down gene expressi	on.		
Form	Lyop	hilized powder			
Gene Symbol	DDX5	DDX55			
Alternative Names KIAA1595; ATP-dependent RNA helicase DDX55; DEAD box protein 55			protein 55		
Entrez Gene 57696 (Human)					
SwissProt Q8NHQ9 (Human)					
Purity > 97%					
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to			h trityl analysis to ensure		
	appro	opriate coupling effici	ency. The oligo is subsequently p	urified by affinity-solid	
	phas	e extraction. The anne	ealed RNA duplex is further analy	zed by mass	
	spect	trometry to verify the	exact composition of the duplex.	Each lot is compared to	
	the p	previous lot by mass sp	pectrometry to ensure maximum	lot-to-lot consistency.	
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	an DDX55 gene. Each	vial contains 5 nmol of lyophilized	siRNA. The duplexes	
	can b	e transfected individu	ally or pooled together to achiev	e knockdown of the	
	targe	target gene, which is most commonly assessed by qPCR or western blot.			
	Con	nponent	15 nmol	30 nmol	
	DDX	(55 siRNA (Human) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

DDX55 siRNA (Human) - B

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

DDX55 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μΙ	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com