

Product Data Sheet

ACE2 siRNA (Human)

Catalog #	Source	Reactivity	Applications			
CRJ1761	Synthetic	н	RNAi			
Description	siRNA	to inhibit ACE2 expr	ession using RNA interference			
Specificity	ACE2	ACE2 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	knock down gene expression.				
Form	Lyoph	Lyophilized powder				
Gene Symbol	ACE2	ACE2				
Alternative N	ames Angio	Angiotensin-converting enzyme 2; ACE-related carboxypeptidase;				
	Angio	tensin-converting en	zyme homolog; ACEH; Metallopro	tease MPROT15		
Entrez Gene	59272	59272 (Human)				
SwissProt	Q9BYI	Q9BYF1 (Human)				
Purity	> 97%	> 97%				
Quality Control Oligonucleotide synthesis is monitored base by base through trity			n trityl analysis to ensure			
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-so				
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass				
	specti	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pr	revious lot by mass s	pectrometry to ensure maximum l	ot-to-lot consistency.		
Components	We of	offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	huma	human ACE2 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can				
	be tra	be transfected individually or pooled together to achieve knockdown of the target				
	gene,	gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	ACE2	2 siRNA (Human) - A	5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

ACE2 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
ACE2 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com