

Product Data Sheet

TSPYL2 siRNA (Human)

Source	Reactivity	Applications		
Synthetic	н	RNAi		
Description siRNA to inhibit TSPYL2 expression using RNA interference				
TSPYL	2 siRNA (Human) is a	target-specific 19-23 nt siRNA oligo duplexes designed to		
knock	down gene expressio	on.		
Lyophi	ilized powder			
TSPYL	TSPYL2			
es CDA1;	DENTT; TSPX; Testis-	specific Y-encoded-like protein 2; TSPY-like protein 2; Cell		
divisio	n autoantigen 1; Cut	aneous T-cell lymphoma-associated antigen se20-4;		
CTCL-a	associated antigen se	20-4; Differentially-expressed nucleolar TGF-beta1 target		
protei	n; Nuclear protein of	79 kDa; NP79		
64061	64061 (Human)			
Q9H20	Q9H2G4 (Human)			
> 97%				
Oligon	ucleotide synthesis i	s monitored base by base through trityl analysis to ensure		
appro	priate coupling efficie	ency. The oligo is subsequently purified by affinity-solid		
phase	extraction. The anne	aled RNA duplex is further analyzed by mass		
spectr	ometry to verify the	exact composition of the duplex. Each lot is compared to		
the pr	evious lot by mass sp	ectrometry to ensure maximum lot-to-lot consistency.		
We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
humar	n TSPYL2 gene. Each	vial contains 5 nmol of lyophilized siRNA. The duplexes		
can be	e transfected individu	ally or pooled together to achieve knockdown of the		
target	gene, which is most	commonly assessed by qPCR or western blot.		
	ynthetic siRNA TSPYL: knock Lyophi TSPYL: es CDA1; divisio CTCL-a protei 64061 Q9H20 > 97% Oligon approj phase spectr the pr We off human can be	ynthetic H siRNA to inhibit TSPYL2 exp TSPYL2 siRNA (Human) is a knock down gene expressio Lyophilized powder TSPYL2 es CDA1; DENTT; TSPX; Testis- division autoantigen 1; Cut CTCL-associated antigen se protein; Nuclear protein of 64061 (Human) Q9H2G4 (Human) > 97% Oligonucleotide synthesis i appropriate coupling efficie phase extraction. The anne spectrometry to verify the the previous lot by mass sp We offers pre-designed set human TSPYL2 gene. Each can be transfected individu		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

Component	15 nmol	30 nmol
TSPYL2 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
TSPYL2 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
TSPYL2 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 µl
6 woll	2 ml	100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

 WEB
 ORDER
 SUPPORT
 CUSTOM

 www.cohesionbio.com
 order@cohesionbio.com
 techsupport@cohesionbio.com
 custom@cohesionbio.com