

Product Data Sheet

ZC3H12A siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRJ2773	Synthetic	н	RNAi		
Description	siRNA	to inhibit ZC3H12A	expression using RNA interferer	nce	
Specificity	ZC3H	ZC3H12A siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kno	ock down gene expre	ssion.		
Form	Lyoph	nilized powder			
Gene Symbol	ZC3H	ZC3H12A			
Alternative N	ames MCPI	MCPIP; MCPIP1; Ribonuclease ZC3H12A; MCP-induced protein 1; Zinc finger CCCH			
	doma	in-containing proteir	12A		
Entrez Gene	80149	9 (Human)			
SwissProt	Q5D1	Q5D1E8 (Human)			
Purity	> 97%	> 97%			
Quality Contr	ol Oligo	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	opriate coupling effici	ency. The oligo is subsequently	purified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spect	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the p	revious lot by mass s	pectrometry to ensure maximu	m lot-to-lot consistency.	
Components	We o	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	an ZC3H12A gene. Ead	ch vial contains 5 nmol of lyoph	ilized siRNA. The duplexes	
	can b	e transfected individ	ually or pooled together to achi	eve knockdown of the	
	targe	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	ZC3ł	H12A siRNA (Human)	- A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

ZC3H12A siRNA (Human) - B	5 nmol x 1	5 nmol x 2
ZC3H12A siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μl
_		10 nM	0.25 μl	1 µl
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 μl
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com