

# **Product Data Sheet**

## EDC3 siRNA (Human)

| Catalog #       | Source     | Reactivity  | Applications                       |                          |  |
|-----------------|------------|---|------------------------------------|--------------------------|--|
| CRJ2776         | Synthetic  | Н   | RNAi                               |                          |  |
| Description     | siRNA      | to inhibit EDC3 expre   | ssion using RNA interference       |                          |  |
| Specificity     | EDC3       | siRNA (Human) is a ta   | rget-specific 19-23 nt siRNA oligo | duplexes designed to     |  |
|                 | knock      | down gene expressio   | n.                                 |                          |  |
| Form            | Lyoph      | ilized powder   |                                    |                          |  |
| Gene Symbol     | EDC3       | EDC3  |                                    |                          |  |
| Alternative N   | ames LSM16 | LSM16; YJDC; YJEFN2; Enhancer of mRNA-decapping protein 3; LSM16 homolog; YjeF        |                                    |                          |  |
|                 | N-terr     | N-terminal domain-containing protein 2; YjeF_N2; hYjeF_N2; YjeF                       |                                    |                          |  |
|                 | domai      | in-containing protein   | 1                                  |                          |  |
| Entrez Gene     | 80153      | (Human)   |                                    |                          |  |
| SwissProt       | Q96F8      | Q96F86 (Human)  |                                    |                          |  |
| Purity          | > 97%      | > 97%   |                                    |                          |  |
| Quality Control | ol Oligor  | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure |                                    |                          |  |
|                 | appro      | priate coupling efficie   | ncy. The oligo is subsequently pu  | rified by affinity-solid |  |
|                 | phase      | extraction. The annea   | aled RNA duplex is further analyz  | ed by mass               |  |
|                 | spectr     | ometry to verify the e  | exact composition of the duplex.   | Each lot is compared to  |  |
|                 | the pr     | evious lot by mass sp   | ectrometry to ensure maximum l     | ot-to-lot consistency.   |  |
| Components      | We of      | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of    |                                    |                          |  |
|                 | huma       | n EDC3 gene. Each via   | l contains 5 nmol of lyophilized s | iRNA. The duplexes can   |  |
|                 | be tra     | nsfected individually o   | or pooled together to achieve kno  | ockdown of the target    |  |
|                 | gene,      | gene, which is most commonly assessed by qPCR or western blot.                        |                                    |                          |  |
|                 | Com        | ponent  | 15 nmol                            | 30 nmol                  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



# **Product Data Sheet**

| EDC3 siRNA (Human) - A | 5 nmol x 1   | 5 nmol x 2   |
|------------------------|--------------|--------------|
| EDC3 siRNA (Human) - B | 5 nmol x 1   | 5 nmol x 2   |
| EDC3 siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control       | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water             | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate     | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|-----------|--------------|---------------------|---------------|--------------|
|           | of medium    | of siRNA            |               | 2000         |
|           |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well   | 100 µl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|           |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|           |              | 100 nM              | 2.5 μl        | 1 μΙ         |
| 24-well   | 500 μl       | 50 nM               | 1.25 μl       | 1 μΙ         |
|           |              | 10 nM               | 0.25 μl       | 1 μl         |
|           |              | 100 nM              | 5 µl          | 2 μΙ         |
| 12-well   | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|           |              | 10 nM               | 0.5 μl        | 2 μΙ         |
|           |              | 100 nM              | 10 µl         | 5 µl         |
| 6-well    | 2 ml         | 50 nM               | 5 μl          | 5 μl         |
| . <u></u> |              | 10 nM               | 1 μl          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | СИЗТОМ                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

## **Product Data Sheet**

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com