

# **Product Data Sheet**

## INO80B siRNA (Human)

| Catalog #     | Source    | Reactivity  | Applications                        |                          |  |  |
|---------------|-----------|---|-------------------------------------|--------------------------|--|--|
| CRJ3098       | Synthetic | н   | RNAi                                |                          |  |  |
| Description   | siRNA     | to inhibit INO80B ex  | pression using RNA interference     |                          |  |  |
| Specificity   | INO80     | INO80B siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to   |                                     |                          |  |  |
|               | knock     | down gene expression  | on.                                 |                          |  |  |
| Form          | Lyoph     | ilized powder   |                                     |                          |  |  |
| Gene Symbol   | INO80     | INO80B  |                                     |                          |  |  |
| Alternative N | ames HMGA | HMGA1L4; PAPA1; ZNHIT4; INO80 complex subunit B; High mobility group AT-hook          |                                     |                          |  |  |
|               | 1-like    | 4; IES2 homolog; hIe  | s2; PAP-1-associated protein 1; PA  | PA-1; Zinc finger HIT    |  |  |
|               | doma      | in-containing protein   | 4                                   |                          |  |  |
| Entrez Gene   | 83444     | 83444 (Human)   |                                     |                          |  |  |
| SwissProt     | Q9C08     | Q9C086 (Human)  |                                     |                          |  |  |
| Purity        | > 97%     | > 97%   |                                     |                          |  |  |
| Quality Contr | ol Oligor | Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure |                                     |                          |  |  |
|               | appro     | priate coupling efficie   | ency. The oligo is subsequently pu  | rified by affinity-solid |  |  |
|               | phase     | extraction. The anne  | ealed RNA duplex is further analyze | ed by mass               |  |  |
|               | spectr    | rometry to verify the   | exact composition of the duplex. I  | Each lot is compared to  |  |  |
|               | the pr    | revious lot by mass sp  | pectrometry to ensure maximum lo    | ot-to-lot consistency.   |  |  |
| Components    | We of     | We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of    |                                     |                          |  |  |
|               | huma      | n INO80B gene. Each   | vial contains 5 nmol of lyophilized | l siRNA. The duplexes    |  |  |
|               | can be    | e transfected individu  | ally or pooled together to achieve  | knockdown of the         |  |  |
|               | target    | target gene, which is most commonly assessed by qPCR or western blot.                 |                                     |                          |  |  |
|               | Com       | ponent  | 15 nmol                             | 30 nmol                  |  |  |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



# **Product Data Sheet**

| INO80B siRNA (Human) - A | 5 nmol x 1   | 5 nmol x 2   |
|--------------------------|--------------|--------------|
| INO80B siRNA (Human) - B | 5 nmol x 1   | 5 nmol x 2   |
| INO80B siRNA (Human) - C | 5 nmol x 1   | 5 nmol x 2   |
| Negative Control         | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water               | 1 ml x 1     | 1 ml x 2     |

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

| Plate   | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
|         | of medium    | of siRNA            |               | 2000         |
|         |              | 100 nM              | 0.5 μl        | 0.25 μl      |
| 96-well | 100 µl       | 50 nM               | 0.25 μl       | 0.25 μl      |
|         |              | 10 nM               | 0.05 μl       | 0.25 μl      |
|         |              | 100 nM              | 2.5 μl        | 1 μΙ         |
| 24-well | 500 μl       | 50 nM               | 1.25 μl       | 1 μl         |
|         |              | 10 nM               | 0.25 μl       | 1 μΙ         |
|         |              | 100 nM              | 5 μl          | 2 μΙ         |
| 12-well | 1 ml         | 50 nM               | 2.5 μl        | 2 μΙ         |
|         |              | 10 nM               | 0.5 μl        | 2 μΙ         |
|         |              | 100 nM              | 10 µl         | 5 µl         |
| 6-well  | 2 ml         | 50 nM               | 5 μl          | 5 μl         |
|         |              | 10 nM               | 1 μl          | 5 μΙ         |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

| WEB                 | ORDER                 | SUPPORT                     | CUSTOM                 |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

## **Product Data Sheet**

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com