

# **Product Data Sheet**

## L3MBTL2 siRNA (Human)

Catalog #	Source	Reactivity	Applicati	ons	
CRJ3189	Synthetic	н	RNAi		
Description	siRNA	to inhibit L3MBTL2 e	expression using RNA interf	ference	
Specificity	L3MB	STL2 siRNA (Human) is	a target-specific 19-23 nt	siRNA oligo duplexes designed	
	to kno	ock down gene expre	ssion.		
Form	Lyoph	ilized powder			
Gene Symbol	L3MB	L3MBTL2			
Alternative N	ames Letha	Lethal(3)malignant brain tumor-like protein 2; H-l(3)mbt-like protein 2; L(3)mbt-like			
	prote	in 2			
Entrez Gene	83746	6 (Human)			
SwissProt	Q969	Q969R5 (Human)			
Purity	> 97%	> 97%			
Quality Contr	ol Oligoi	nucleotide synthesis i	s monitored base by base	through trityl analysis to ensure	
	appro	priate coupling effici	ency. The oligo is subseque	ently purified by affinity-solid	
	phase	e extraction. The anne	ealed RNA duplex is further	r analyzed by mass	
	spect	rometry to verify the	exact composition of the c	luplex. Each lot is compared to	
	the p	revious lot by mass sp	pectrometry to ensure max	imum lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	n L3MBTL2 gene. Eac	h vial contains 5 nmol of ly	ophilized siRNA. The duplexes	
	can b	e transfected individu	ally or pooled together to	achieve knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	L3M	BTL2 siRNA (Human)	- A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

#### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

L3MBTL2 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
L3MBTL2 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
_		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
_		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 µl	5 µl
		10 nM	1 μΙ	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com