

Product Data Sheet

PHYHIPL siRNA (Human)

Catalog #	Source	Reactivity	Application	ons	
CRJ3486	Synthetic	н	RNAi		
Description	siRNA	to inhibit PHYHIPL ex	pression using RNA interfe	erence	
Specificity	РНҮН	PHYHIPL siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kno	ock down gene expres	ssion.		
Form	Lyoph	ilized powder			
Gene Symbol	РНҮН	PHYHIPL			
Alternative N	ames KIAA1	KIAA1796; Phytanoyl-CoA hydroxylase-interacting protein-like			
Entrez Gene 84457 (Human)					
SwissProt Q96FC7 (Human)					
Purity > 97%					
Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis to			through trityl analysis to ensure		
	appro	priate coupling efficie	ency. The oligo is subseque	ently purified by affinity-solid	
	phase	extraction. The anne	aled RNA duplex is further	analyzed by mass	
	specti	rometry to verify the	exact composition of the c	luplex. Each lot is compared to	
	the pr	revious lot by mass sp	ectrometry to ensure max	imum lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	human PHYHIPL gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes			
	can be	e transfected individu	ally or pooled together to	achieve knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	PHY	HIPL siRNA (Human) -	A 5 nmol x 1	5 nmol x 2	
	51.54			-	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-

5 nmol x 1

5 nmol x 2

PHYHIPL siRNA (Human) - B

Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

DEPC Water	1 ml x 1	1 ml x 2
Negative control	2.5 111101 X 1	2.5 111101 X 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
PHYHIPL siRNA (Human) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μΙ	5 µl
		10 nM	1 µl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com