

Product Data Sheet

USP38 siRNA (Human)

Catalog #	Source	Reactivity	Applications			
CRJ3550	Synthetic	н	RNAi			
Description	siRNA	to inhibit USP38 exp	ression using RNA interference	e		
Specificity	USP38	USP38 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
	knock	down gene expression	on.			
Form	Lyoph	ilized powder				
Gene Symbol	USP38	USP38				
Alternative N	ames KIAA1	KIAA1891; Ubiquitin carboxyl-terminal hydrolase 38; Deubiquitinating enzyme 38;				
	HP43.	8KD; Ubiquitin thioe	sterase 38; Ubiquitin-specific-r	processing protease 38		
Entrez Gene	84640) (Human)				
SwissProt	Q8NB	Q8NB14 (Human)				
Purity	> 97%	> 97%				
Quality Control Oligonucleotide synthesis			s monitored base by base thro	ough trityl analysis to ensure		
	appro	priate coupling efficient	ency. The oligo is subsequently	y purified by affinity-solid		
	phase	extraction. The anne	ealed RNA duplex is further an	alyzed by mass		
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to				
	the pr	evious lot by mass sp	pectrometry to ensure maximu	ım lot-to-lot consistency.		
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of				
	huma	human USP38 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can				
	be tra	be transfected individually or pooled together to achieve knockdown of the target				
	gene,	gene, which is most commonly assessed by qPCR or western blot.				
	Com	ponent	15 nmol	30 nmol		
	USP3	38 siRNA (Human) - A	5 nmol x 1	5 nmol x 2		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

USP38 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
USP38 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
_		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 µl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com