

Product Data Sheet

KDM2B siRNA (Human)

| Catalog # | Source | Reactivity | Applications |
|----------------|-----------|-------------------------|--|
| CRJ3575 | Synthetic | н | RNAi |
| Description | siR | NA to inhibit KDM2B | expression using RNA interference |
| Specificity | KD | M2B siRNA (Human) | is a target-specific 19-23 nt siRNA oligo duplexes designed to |
| | kno | ock down gene expre | ssion. |
| Form | Lyc | ophilized powder | |
| Gene Symbol | KD | M2B | |
| Alternative Na | ames CX | XC2; FBL10; FBXL10; J | HDM1B; PCCX2; Lysine-specific demethylase 2B; CXXC-type |
| | zin | c finger protein 2; F-b | ox and leucine-rich repeat protein 10; F-box protein FBL10; |
| | F-b | oox/LRR-repeat protei | n 10; JmjC domain-containing histone demethylation |
| | pro | otein 1B; Jumonji dom | nain-containing EMSY-interactor methyltransferase motif |
| | pro | otein; Protein JEMMA | ; Protein-containing CXXC domain 2; [Histone-H3]-lysine-36 |
| | dei | methylase 1B | |
| Entrez Gene | 846 | 678 (Human) | |
| SwissProt | Q8 | NHM5 (Human) | |
| Purity | > 9 | 7% | |
| Quality Contro | ol Oli | gonucleotide synthes | is is monitored base by base through trityl analysis to ensure |
| | ар | propriate coupling eff | iciency. The oligo is subsequently purified by affinity-solid |
| | pha | ase extraction. The ar | nealed RNA duplex is further analyzed by mass |
| | spe | ectrometry to verify t | ne exact composition of the duplex. Each lot is compared to |
| | the | e previous lot by mass | spectrometry to ensure maximum lot-to-lot consistency. |
| Components | We | e offers pre-designed | sets of 3 different target-specific siRNA oligo duplexes of |
| | hu | man KDM2B gene. Ea | ch vial contains 5 nmol of lyophilized siRNA. The duplexes |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

can be transfected individually or pooled together to achieve knockdown of the

target gene, which is most commonly assessed by qPCR or western blot.

| Component | 15 nmol | 30 nmol |
|-------------------------|--------------|--------------|
| KDM2B siRNA (Human) - A | 5 nmol x 1 | 5 nmol x 2 |
| KDM2B siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
| KDM2B siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μl of DEPC water to get a final concentration of 20 μM.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 μl | 2 μl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μl |
| | | 10 nM | 0.5 μl | 2 µl |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



For research purposes only, not for human use

Product Data Sheet

| | | 100 nM | 10 µl | 5 μl |
|--------|------|--------|-------|------|
| 6-well | 2 ml | 50 nM | 5 µl | 5 µl |
| | | 10 nM | 1 µl | 5 µl |

Storage/Stability Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com