

# **Product Data Sheet**

## USMG5 siRNA (Human)

rce					
h	Reactivity	Applications			
netic	Н	RNAi			
siRNA	siRNA to inhibit USMG5 expression using RNA interference				
USMO	USMG5 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to				
knock	down gene expressi	on.			
Lyoph	ilized powder	zed powder			
USMO	55				
DAPIT	DAPIT; HCVFTP2; Up-regulated during skeletal muscle growth protein 5;				
Diabe	tes-associated protei	n in insulin-sensitive tissues; HCV F-transactivated protein			
2					
84833	8 (Human)				
SwissProt Q96IX5 (Hu					
> 97%	> 97%				
Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure				
appro	priate coupling effici	ency. The oligo is subsequently purified by affinity-solid			
phase	extraction. The anno	ealed RNA duplex is further analyzed by mass			
spectr	rometry to verify the	exact composition of the duplex. Each lot is compared to			
the pr	revious lot by mass sp	pectrometry to ensure maximum lot-to-lot consistency.			
Components We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of					
human USMG5 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes					
can be	e transfected individu	ally or pooled together to achieve knockdown of the			
target	gene, which is most	commonly assessed by qPCR or western blot.			
Com	ponent	15 nmol 30 nmol			
	USMO knock Lyoph USMO DAPIT Diabe 2 84833 Q96IX > 97% Oligon appro phase spect the pr We of huma can bo target	siRNA to inhibit USMG5 ex USMG5 siRNA (Human) is a knock down gene expression Lyophilized powder USMG5 DAPIT; HCVFTP2; Up-regular Diabetes-associated protein 2 84833 (Human) Q96IX5 (Human) > 97% Oligonucleotide synthesis in appropriate coupling efficient phase extraction. The annea spectrometry to verify the the previous lot by mass sp We offers pre-designed set human USMG5 gene. Each can be transfected individu			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



# **Product Data Sheet**

USMG5 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
USMG5 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
USMG5 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μΙ
24-well	500 μl	50 nM	1.25 μl	1 μl
		10 nM	0.25 μl	1 μΙ
		100 nM	5 µl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 µl	5 µl
		10 nM	1 μl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

## **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



For research purposes only, not for human use

## **Product Data Sheet**

**Storage/Stability** Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com