

Product Data Sheet

KRTAP4-3 siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRJ3768	Synthetic	н	RNAi		
Description	siRNA	to inhibit KRTAP4-3	expression using RNA interfere	nce	
Specificity	KRTAF	KRTAP4-3 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed			
	to kno	ock down gene expre	ssion.		
Form	Lyoph	ilized powder			
Gene Symbol	KRTAF	KRTAP4-3			
Alternative N	ames KAP4.	KAP4.3; KRTAP4.3; Keratin-associated protein 4-3; Keratin-associated protein 4.3;			
	Ultrah	nigh sulfur keratin-as	sociated protein 4.3		
Entrez Gene	85290) (Human)			
SwissProt	Q9BYI	Q9BYR4 (Human)			
Purity	> 97%	> 97%			
Quality Contr	ol Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	priate coupling effici	ency. The oligo is subsequently	purified by affinity-solid	
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the pr	evious lot by mass sp	pectrometry to ensure maximu	m lot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	n KRTAP4-3 gene. Ea	ch vial contains 5 nmol of lyoph	ilized siRNA. The duplexes	
	can be	e transfected individu	ally or pooled together to achi	eve knockdown of the	
	target	target gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	KRTA	P4-3 siRNA (Human)	- A 5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

KRTAP4-3 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
KRTAP4-3 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
		100 nM	10 µl	5 μΙ
6-well	2 ml	50 nM	5 μl	5 μΙ
		10 nM	1 μl	5 μΙ

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com