

Product Data Sheet

EAF1 siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRJ3797	Synthetic	Н	RNAi		
Description	siRNA	to inhibit EAF1 expres	ssion using RNA interference		
Specificity	EAF1	EAF1 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	down gene expressio	n.		
Form	Lyoph	Lyophilized powder			
Gene Symbol	EAF1	EAF1			
Alternative Na	ames ELL-as	ELL-associated factor 1			
Entrez Gene	85403	85403 (Human)			
SwissProt	Q96JC	Q96JC9 (Human)			
Purity	> 97%	> 97%			
Quality Contro	ol Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
	appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid			
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the pr	evious lot by mass spe	ectrometry to ensure maximum l	ot-to-lot consistency.	
Components	We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
	huma	human EAF1 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can			
	be tra	be transfected individually or pooled together to achieve knockdown of the target			
	gene,	gene, which is most commonly assessed by qPCR or western blot.			
	Com	ponent	15 nmol	30 nmol	
	EAF1	siRNA (Human) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

EAF1 siRNA (Human) - B

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

DEPC Water		1 ml x 1	1 ml x 2
Negative Contro	bl	2.5 nmol x 1	2.5 nmol x 2
EAF1 siRNA (Hu	man) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 µl
6-well	2 ml	50 nM	5 μl	5 µl
		10 nM	1 µl	5 μl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com