

Product Data Sheet

SHF siRNA (Human)

Catalog #	Source	Reactivity	Applications		
CRJ3957	Synthetic	н	RNAi		
Description	siRNA	siRNA to inhibit SHF expression using RNA interference			
Specificity	SHF si	SHF siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to			
	knock	knock down gene expression.			
Form	Lyoph	Lyophilized powder			
Gene Symbol	SHF	SHF			
Alternative N	ames SH2 de	SH2 domain-containing adapter protein F			
Entrez Gene	90525	90525 (Human)			
SwissProt	Q7M4	Q7M4L6 (Human)			
Purity	> 97%	> 97%			
Quality ControlOligonucleotide synthesis is monitored base by base thro			monitored base by base throug	h trityl analysis to ensure	
appropriate coupling efficiency. The oligo is s		cy. The oligo is subsequently pu	subsequently purified by affinity-solid		
	phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
	spectr	spectrometry to verify the exact composition of the duplex. Each lot is compared to			
	the pr	evious lot by mass spe	ctrometry to ensure maximum	lot-to-lot consistency.	
Components We offers pre-designed sets of 3 different target-specific siRNA oligo du			NA oligo duplexes of		
	humai	human SHF gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can			
be transfected individually or pooled together to achieve knockdown			ockdown of the target		
	gene, which is most commonly assessed by qPCR or western blot.			n blot.	
	Com	ponent	15 nmol	30 nmol	
	SHF s	siRNA (Human) - A	5 nmol x 1	5 nmol x 2	

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

5 nmol x 1

5 nmol x 2

SHF siRNA (Human) - B

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



Product Data Sheet

	DEPC Water	1 ml x 1	1 ml x 2
r		1 ml v 1	1 ml v 2
1	Negative Control	2.5 nmol x 1	2.5 nmol x 2
<u>s</u>	6HF siRNA (Human) - C	5 nmol x 1	5 nmol x 2

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 µl
		10 nM	0.25 μl	1 µl
		100 nM	5 µl	2 µl
12-well	1 ml	50 nM	2.5 μl	2 µl
		10 nM	0.5 μl	2 µl
		100 nM	10 µl	5 μl
6-well	2 ml	50 nM	5 µl	5 µl
		10 nM	1 µl	5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com