

# **Product Data Sheet**

### AP5B1 siRNA (Human)

Reactivity	Applications				
tic H	RNAi				
escription siRNA to inhibit AP5B1 expression using RNA interference					
AP5B1 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to					
knock down gene expression.					
Lyophilized powder	hilized powder				
AP5B1					
Names AP-5 complex subunit beta-1; Adapter-related protein complex 5 beta subunit;					
Beta5					
91056 (Human)					
Q2VPB7 (Human)					
> 97%					
Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure					
appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid					
phase extraction. The annealed RNA duplex is further analyzed by mass					
<ul> <li>spectrometry to verify the exact composition of the duplex. Each lot is compared to the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency.</li> <li>We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of human AP5B1 gene. Each vial contains 5 nmol of lyophilized siRNA. The duplexes can</li> </ul>					
			be transfected individually or pooled t	ogether to achieve know	kdown of the target
			gene, which is most commonly assessed by qPCR or western blot.		
			Component	15 nmol	30 nmol
	tic H siRNA to inhibit AP5B1 expression usin AP5B1 siRNA (Human) is a target-speci knock down gene expression. Lyophilized powder AP5B1 AP-5 complex subunit beta-1; Adapter Beta5 91056 (Human) Q2VPB7 (Human) > 97% Oligonucleotide synthesis is monitored appropriate coupling efficiency. The ol phase extraction. The annealed RNA d spectrometry to verify the exact comp the previous lot by mass spectrometry We offers pre-designed sets of 3 differ human AP5B1 gene. Each vial contains be transfected individually or pooled t gene, which is most commonly assessed	tic H RNAi siRNA to inhibit AP5B1 expression using RNA interference AP5B1 siRNA (Human) is a target-specific 19-23 nt siRNA oligo knock down gene expression. Lyophilized powder AP5B1 AP-5 complex subunit beta-1; Adapter-related protein comple Beta5 91056 (Human) Q2VPB7 (Human) > 97% Oligonucleotide synthesis is monitored base by base through appropriate coupling efficiency. The oligo is subsequently puri phase extraction. The annealed RNA duplex is further analyze spectrometry to verify the exact composition of the duplex. Ea the previous lot by mass spectrometry to ensure maximum lo We offers pre-designed sets of 3 different target-specific siRN, human AP5B1 gene. Each vial contains 5 nmol of lyophilized si be transfected individually or pooled together to achieve know gene, which is most commonly assessed by qPCR or western b			

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



# **Product Data Sheet**

AP5B1 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
AP5B1 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

**Directions for Use** 

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 μl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 μl	2 μl
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well		100 nM	10 µl	5 µl
	2 ml	50 nM	5 µl	5 μΙ
		10 nM	1 μΙ	5 μΙ

#### Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com