

Product Data Sheet

KLHL29 siRNA (Human)

| Catalog # | Source | Reactivity | Applications | | | |
|---|---|---|----------------------------------|----------------------------|--|--|
| CRJ4390 | Synthetic | н | RNAi | | | |
| Description | Description siRNA to inhibit KLHL29 expression using RNA interference | | | е | | |
| Specificity | KLHL2 | KLHL29 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed to | | | | |
| | knock | knock down gene expression. | | | | |
| Form | Lyoph | Lyophilized powder | | | | |
| Gene Symbol | KLHL2 | KLHL29 | | | | |
| Alternative N | ames KBTBE | KBTBD9; KIAA1921; Kelch-like protein 29; Kelch repeat and BTB domain-containing | | | | |
| | protei | in 9 | | | | |
| Entrez Gene | 11481 | 114818 (Human) | | | | |
| SwissProt | Q96C | Q96CT2 (Human) | | | | |
| Purity | > 97% | > 97% | | | | |
| Quality Control Oligonucleotide synthesis is monitored base by base through trityl analysis | | | ugh trityl analysis to ensure | | | |
| | appropriate coupling efficiency. The oligo is subsequently purified by affinity-s | | | purified by affinity-solid | | |
| | phase | phase extraction. The annealed RNA duplex is further analyzed by mass | | | | |
| | spectr | spectrometry to verify the exact composition of the duplex. Each lot is compared to | | | | |
| | the pr | the previous lot by mass spectrometry to ensure maximum lot-to-lot consistency. | | | | |
| Components We offers pre-designed sets of 3 different target-specific siRNA oligo du | | | iRNA oligo duplexes of | | | |
| | huma | n KLHL29 gene. Each | vial contains 5 nmol of lyophili | zed siRNA. The duplexes | | |
| | can be | e transfected individu | ally or pooled together to achi | eve knockdown of the | | |
| | target | target gene, which is most commonly assessed by qPCR or western blot. | | | | |
| | Com | ponent | 15 nmol | 30 nmol | | |
| | KLHL | .29 siRNA (Human) - / | 4 5 nmol x 1 | 5 nmol x 2 | | |

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |



Product Data Sheet

| KLHL29 siRNA (Human) - B | 5 nmol x 1 | 5 nmol x 2 |
|--------------------------|--------------|--------------|
| KLHL29 siRNA (Human) - C | 5 nmol x 1 | 5 nmol x 2 |
| Negative Control | 2.5 nmol x 1 | 2.5 nmol x 2 |
| DEPC Water | 1 ml x 1 | 1 ml x 2 |

Directions for Use

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250 μ l of DEPC water to get a final concentration of 20 μ M.

| Plate | Final volume | Final concentration | siRNA (20 μM) | Lipofectamin |
|---------|--------------|---------------------|---------------|--------------|
| | of medium | of siRNA | | 2000 |
| | | 100 nM | 0.5 μl | 0.25 μl |
| 96-well | 100 µl | 50 nM | 0.25 μl | 0.25 μl |
| _ | | 10 nM | 0.05 μl | 0.25 μl |
| | | 100 nM | 2.5 μl | 1 μl |
| 24-well | 500 μl | 50 nM | 1.25 μl | 1 μΙ |
| _ | | 10 nM | 0.25 μl | 1 μΙ |
| | | 100 nM | 5 μl | 2 μl |
| 12-well | 1 ml | 50 nM | 2.5 μl | 2 μΙ |
| _ | | 10 nM | 0.5 μl | 2 μΙ |
| | | 100 nM | 10 µl | 5 µl |
| 6-well | 2 ml | 50 nM | 5 μl | 5 μΙ |
| | | 10 nM | 1 μl | 5 μΙ |

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

COHESION BIOSCIENCES LIMITED

| WEB | ORDER | SUPPORT | CUSTOM |
|---------------------|-----------------------|-----------------------------|------------------------|
| www.cohesionbio.com | order@cohesionbio.com | techsupport@cohesionbio.com | custom@cohesionbio.com |