

# **Product Data Sheet**

## IL22RA2 siRNA (Human)

Source	Reactivity	Applications		
Synthetic	Н	RNAi		
siRNA	to inhibit IL22RA2 ex	pression using RNA interference		
IL22R/	IL22RA2 siRNA (Human) is a target-specific 19-23 nt siRNA oligo duplexes designed			
to kno	ock down gene expres	ssion.		
Lyoph	ilized powder			
IL22R/	42			
nes Interle	Interleukin-22 receptor subunit alpha-2; IL-22 receptor subunit alpha-2;			
IL-22R	IL-22R-alpha-2; IL-22RA2; Cytokine receptor class-II member 10; Cytokine receptor			
family	10; Cytokine receptor family type 2. soluble 1; CRF2-S1;			
Interle	eukin-22-bin			
11637	116379 (Human)			
Q969J	Q969J5 (Human)			
> 97%	> 97%			
Oligor	Oligonucleotide synthesis is monitored base by base through trityl analysis to ensure			
appro	appropriate coupling efficiency. The oligo is subsequently purified by affinity-solid			
phase	phase extraction. The annealed RNA duplex is further analyzed by mass			
spectr	rometry to verify the	exact composition of the duplex. Each lot is compared to		
the pr	evious lot by mass sp	ectrometry to ensure maximum lot-to-lot consistency.		
We of	We offers pre-designed sets of 3 different target-specific siRNA oligo duplexes of			
huma	n IL22RA2 gene. Each	vial contains 5 nmol of lyophilized siRNA. The duplexes		
can be	e transfected individu	ally or pooled together to achieve knockdown of the		
target	gene, which is most	commonly assessed by qPCR or western blot.		
	Synthetic siRNA IL22R/ to kno Lyoph IL22R/ I	Synthetic H   siRNA to inhibit IL22RA2 ex   IL22RA2 siRNA (Human) is a   to knock down gene express   Lyophilized powder   IL22RA2   Interleukin-22 receptor sub   IL-22R-alpha-2; IL-22RA2; C   family 2 member 10; CRF2-   Interleukin-22-bin   116379 (Human)   Q969J5 (Human)   > 97%   Oligonucleotide synthesis is   appropriate coupling efficies   phase extraction. The anne   spectrometry to verify the   the previous lot by mass sp   We offers pre-designed set   human IL22RA2 gene. Each   can be transfected individu		

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

Component	15 nmol	30 nmol
IL22RA2 siRNA (Human) - A	5 nmol x 1	5 nmol x 2
IL22RA2 siRNA (Human) - B	5 nmol x 1	5 nmol x 2
IL22RA2 siRNA (Human) - C	5 nmol x 1	5 nmol x 2
Negative Control	2.5 nmol x 1	2.5 nmol x 2
DEPC Water	1 ml x 1	1 ml x 2

#### **Directions for Use**

We recommends transfection with 10 nM - 100 nM siRNA 48 to 72 hours prior to cell lysis. Before resuspending, briefly centrifuge the tube to ensure the lyophilized siRNA is at the bottom of the tube. Resuspend the siRNA oligos to an appropriate concentration with DEPC water. For example, resuspend one tube of 5 nmol siRNA oligo in 250  $\mu$ l of DEPC water to get a final concentration of 20  $\mu$ M.

Plate	Final volume	Final concentration	siRNA (20 μM)	Lipofectamin
	of medium	of siRNA		2000
		100 nM	0.5 μl	0.25 μl
96-well	100 µl	50 nM	0.25 μl	0.25 μl
		10 nM	0.05 μl	0.25 μl
		100 nM	2.5 μl	1 µl
24-well	500 μl	50 nM	1.25 μl	1 μΙ
		10 nM	0.25 μl	1 μΙ
		100 nM	5 µl	2 μΙ
12-well	1 ml	50 nM	2.5 μl	2 μΙ
		10 nM	0.5 μl	2 μΙ
6-well	2 ml	100 nM	10 µl	5 µl
o-well	2 ml	50 nM	5 µl	5 μΙ

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEB	ORDER	SUPPORT	CUSTOM
www.cohesionbio.com	order@cohesionbio.com	techsupport@cohesionbio.com	custom@cohesionbio.com



## **Product Data Sheet**

10 nM

1 µl

5 µl

Storage/Stability

Shipped at 4 °C. Store at -20 °C for one year.

Application key: E- ELISA, WB- Western blot, IH- Immunohistochemistry, IF- Immunofluorescence, FC- Flow cytometry, IC-Immunocytochemistry, IP- Immunoprecipitation, ChIP- Chromatin Immunoprecipitation, EMSA- Electrophoretic Mobility Shift Assay, BL- Blocking, SE- Sandwich ELISA, CBE- Cell-based ELISA, RNAi- RNA interference Species reactivity key: H- Human, M- Mouse, R- Rat, B- Bovine, C- Chicken, D- Dog, G- Goat, Mk- Monkey, P- Pig, Rb-Rabbit, S- Sheep, Z- Zebrafish

### **COHESION BIOSCIENCES LIMITED**

WEBORDERSUPPORTCUSTOMwww.cohesionbio.comorder@cohesionbio.comtechsupport@cohesionbio.comcustom@cohesionbio.com